arXiv:0710.4998v2 [math.NT] 4 Sep 2008

BINARY NORMAL REGULAR HERMITIAN LATTICES OVER IMAGINARY QUADRATIC FIELDS

BYEONG MOON KIM, JI YOUNG KIM, AND POO-SUNG PARK

Abstract. We call a positive definite Hermitian lattice regular if it represents all integers, which can be represented locally by the lattice. We investigate binary regular Hermitian lattices over imaginary quadratic fields $\mathbb{Q}(\sqrt{-m})$ and provide a complete list of the (normal) binary regular Hermitian lattices.

1. INTRODUCTION

Mathematicians or number theorists have been pursuing, for a long time, an algorithm to check whether a given Diophantine equation has a solution without solving the equation directly. A basic method to do this is to investigate the behavior of the residues modulo some numbers. This method may show the insolvability of a Diophantine equation. But, it is still considered to be a hard problem to determine when this method guarantees the solvability.

Dickson first called a positive definite quadratic form f *regular* if $f = n$ has an integral solution for each n such that $f \equiv n \pmod{m}$ has solutions for all positive integers m. He computed all regular forms $x^2 + ay^2 + bz^2$, as a generalization of the famous unsolved problem, Euler's idoneal number a admitting $x^2 + ay^2$ to be regular [5]. Jones extended Dickson's results to the form $ax^2 + by^2 + cz^2$ [13]. His work contained a candidate and it was immediately solved by himself and Pall [14].

The outstanding result about regular quadratic forms was achieved by Watson. He showed that there are finitely many equivalence classes of primitive positive definite regular ternary quadratic forms [22], [23]. The complete list of 913 regular ternary forms including 22 candidates was given by Jagy, Kaplansky and Schiemann [12]. On the contrary, Earnest found an infinite family of regular quaternary forms [6] and the first author carried out the determination of all regular diagonal quaternary forms [15].

The regularity of integral quadratic forms is naturally generalized to that of lattices over totally real algebraic number fields. Recently the analogue of Watson's finiteness result for regular positive definite ternary quadratic lattices over the ring $\mathfrak{o}$ of $\mathbb{Q}(\sqrt{5})$ was proved [2].

Regular Hermitian lattices over imaginary quadratic fields are defined in a similar way. If a Hermitian lattice represents all positive integers, it is trivially regular. We call such lattices *universal*. The universal Hermitian lattices were concentrative subjects studied by many mathematicians including the authors in the last couple of decades [7], [10], [16], [17].

The *regular* Hermitian lattices were also investigated and the finiteness of binary *normal* regular Hermitian lattices was proved by Earnest and Khosravani [8]. Besides, several binary regular Hermitian diagonal lattices including a candidate $\langle 1, 14\rangle$ over $\mathbb{Q}(\sqrt{-7})$ were listed by Rokicki [21]. But the inventory was limited to the diagonal lattices, that is, $\mathfrak{a}_1 v_1 \perp \mathfrak{a}_2 v_2$ with two ideals $\mathfrak{a}_1, \mathfrak{a}_2 \subseteq \mathfrak{o}$ and two vectors v_1, v_2. The obstruction against studying Hermitian lattices was that the matrix

---

Date: Received: date / Accepted: date.

2000 *Mathematics Subject Classification.* Primary 11E39; Secondary 11E20, 11E41.

presentation was unprovided. The authors, however, developed the formal matrix presentation and were able to delve into universality and regularity of Hermitian lattices. So we obtained a complete list of positive definite binary normal regular Hermitian lattices including nondiagonal lattices with complete proofs. Rokicki's lattice $\langle 1, 14\rangle$ over $\mathbb{Q}(\sqrt{-7})$ is proved to be regular. To do this, we developed a new method to calculate numbers represented by a quaternary quadratic form. The binary subnormal regular lattices will appear in the next articles.

2. Preliminaries

In this section we give some notations and terminologies, which are adopted from [11] and [19]. Let $\mathfrak{o}$ be the ring of integers of the imaginary quadratic field $E = \mathbb{Q}(\sqrt{-m})$. We have that $\mathfrak{o} = \mathbb{Z}[\omega]$ with $\omega = \sqrt{-m}$ if $m \not\equiv 3 \pmod 4$ and $\omega = \frac{1+\sqrt{-m}}{2}$ if $m \equiv 3 \pmod 4$. A Hermitian space V is a vector space over $\mathbb{Q}(\sqrt{-m})$ with a map $H = H_V : V \times V \to \mathbb{Q}(\sqrt{-m})$ satisfying the following conditions:

1. $H(v, w) = \overline{H(w, v)}$ for $v, w \in V$,
2. $H(v_1 + v_2, w) = H(v_1, w) + H(v_2, w)$ for $v_1, v_2, w \in V$,
3. $H(av, w) = aH(v, w)$ for $a \in \mathbb{Q}(\sqrt{-m})$ and $v, w \in V$.

For brevity, we write $H(v) = H(v, v)$. A Hermitian lattice L is defined as a finitely generated $\mathfrak{o}$-module in the Hermitian space V. From condition (1), we know that

$$H(v) = H(v, v) = \overline{H(v, v)} = \overline{H(v)}.$$

Hence $H(v)$ is always a rational integer. If $a = H(v)$ for some $v \in L$, we say that a is represented by L and denote it by $a \to L$. If a cannot be represented by L, we denote it by $a \not\to L$. Through this article, we assume that L is positive definite, i.e., $H(v) > 0$ for nonzero vectors $v \in L$.

The localization of a lattice L is defined by $L_{\mathfrak{p}} = \mathfrak{o}_{\mathfrak{p}} \otimes_{\mathfrak{o}} L$ and E-ification of L is defined by $EL = E \otimes_{\mathfrak{o}} L$. If $n \to L_{\mathfrak{p}}$ for all primes $\mathfrak{p}$ including ∞, then we write $n \to \operatorname{gen} L$. The regularity of a Hermitian lattice L can be rephrased as follows: if $n \to \operatorname{gen} L$, then $n \to L$. Thus if the class number of L is one, then L is trivially regular.

If a regular lattice L is locally universal over $\mathfrak{o}_{\mathfrak{p}}$ for all prime ideals $\mathfrak{p}$, then L is universal. Since all universal Hermitian lattices are already classified [7], [10], [17], we only consider nonuniversal regular lattices through this article.

We define two ideals related to values of H. The norm $\mathfrak{n}L$ of L is the $\mathfrak{o}$-ideal generated by the set $\{H(v) | v \in L\}$. The scale $\mathfrak{s}L$ of L is the $\mathfrak{o}$-ideal generated by the set $\{H(v, w) | v, w \in L\}$. It is clear that $\mathfrak{n}L \subseteq \mathfrak{s}L$. If $\mathfrak{n}L = \mathfrak{s}L$, then we call L *normal*. Otherwise, we call L *subnormal*. We investigate normal lattices in this article.

The lattice can be written as

$$L = \mathfrak{a}_1 v_1 + \mathfrak{a}_2 v_2 + \cdots + \mathfrak{a}_n v_n$$

with ideals $\mathfrak{a}_i \subset \mathfrak{o}$ and vectors $v_i \in V$. If these vectors are linearly independent over $\mathbb{Q}(\sqrt{-m})$, then we say that L is a n-ary lattice and $\operatorname{rank} L = n$. The significant invariant of L is the *volume* defined as

$$\mathfrak{v}L = (\mathfrak{a}_1\overline{\mathfrak{a}}_1)(\mathfrak{a}_2\overline{\mathfrak{a}}_2)\cdots(\mathfrak{a}_n\overline{\mathfrak{a}}_n)\det(H(v_i, v_j))_{1\le i,j\le n}.$$

Note that this ideal is invariant for equivalent lattices, i.e., $\mathfrak{v}L = \mathfrak{v}(\phi L)$ for an isometry ϕ and the volumes of sublattices are contained in $\mathfrak{v}L$.

If L is a free $\mathfrak{o}$-module, then we can write $L = \mathfrak{o}v_1 + \cdots + \mathfrak{o}v_n$. The matrix $M_L = (H(v_i, v_j))_{1\le i,j\le n}$ is called the Gram matrix of L and is a matrix presentation of L. If the matrix is diagonal, we denote it by $\langle H(v_1), H(v_2), \ldots, H(v_n)\rangle$. But, if

L is not a free $\mathfrak{o}$-module, then $L = \mathfrak{o}v_1 + \cdots + \mathfrak{o}v_{n-1} + \mathfrak{a}v_n$ for some ideal $\mathfrak{a} \subset \mathfrak{o}$ [19, 81:5]. Since any ideal in $\mathfrak{o}$ is generated by at most two elements, we can write $L = \mathfrak{o}v_1 + \cdots + \mathfrak{o}v_{n-1} + (\alpha, \beta)\mathfrak{o}v_n$ for some $\alpha, \beta \in \mathfrak{o}$. Therefore, we consider the following $(n+1) \times (n+1)$-matrix as a formal Gram matrix for L:

$$M_L = \begin{pmatrix} H(v_1, v_1) & \dots & H(v_1, \alpha v_n) & H(v_1, \beta v_n) \\ \vdots & \ddots & \vdots & \vdots \\ H(\alpha v_n, v_1) & \dots & H(\alpha v_n, \alpha v_n) & H(\alpha v_n, \beta v_n) \\ H(\beta v_n, v_1) & \dots & H(\beta v_n, \alpha v_n) & H(\beta v_n, \beta v_n) \end{pmatrix}.$$

Note that this matrix is positive semi-definite, but this represents an n-ary positive definite Hermitian lattice. A scaled lattice L^a obtained from the Hermitian form $H_{L^a} = aH_L$ with $a \in \mathbb{Z}$. If M is a matrix representation of a lattice L, we write aM for the matrix representation of a scaled lattice L^a.

If the (formal) Gram matrix (a_{ij}) is called *Minkowski-reduced* if it satisfies the following conditions:

(1) $a_{ii} \leq a_{jj}$ for $i < j$,

(2) $a_{ii} \leq |2a_{ij,1}|$ and $a_{ii} \leq |2a_{ij,2}|$ for $a_{ij} = a_{ij,1} + \omega a_{ij,2}$ with $i > j$.

A Hermitian lattice is defined over $\mathbb{Z}[\omega]$ and an algebraic integer is of the form $x + y\omega$. So we can define values of $2n$-ary quadratic form $\widetilde{H}(x_1, y_1, \ldots, x_n, y_n)$ over $\mathbb{Z}$ as values of n-ary Hermitian form $H(x_1 + y_1\omega, \ldots, x_n + y_n\omega)$ over $\mathbb{Z}[\omega]$. We call this quadratic form *associated* with the Hermitian lattice [7]. It is sometimes convenient to consider the associated quadratic lattice instead of the Hermitian lattice. To distinguish the associated lattice from the original one, we use the subscript $\mathbb{Z}$. For instance, the quadratic form $\langle 1, 1, 1, 1\rangle_{\mathbb{Z}}$ is associated with the Hermitian form $\langle 1, 1\rangle$ over $\mathbb{Z}[\sqrt{-1}]$. We abuse both terms quadratic *forms* and quadratic *lattices*. So we might say that a quadratic form has a sublattice.

3. Watson transformation

For a positive integer t and a Hermitian lattice L, let

$$\Lambda_t(L) = \{v \in L | H(v, w) \equiv 0 \pmod{t} \text{ for all } w \in L\}.$$

Let the *Watson transformation* $\lambda_t(L)$ of L be the lattice defined by

$$\lambda_t(L) = \Lambda_t(L)^{\frac{1}{a}},$$

where a is the maximal positive integer which divides all elements of $\{B(v, w) | v, w \in \Lambda_t(L)\}$. If L is regular, then $\lambda_t(L)$ is also regular [24], [3].

Throughout the remainder of this article, L will always means an $\mathfrak{o}$-lattice on a positive definite binary Hermitian space over $E = \mathbb{Q}(\sqrt{-m})$, where m is a positive integer. We may assume that all lattices L under discussion are primitive, since if a lattice is regular, then so is any multiple of that lattice. Let p be a rational odd prime and $\mathfrak{p} = \mathfrak{p}(p)$ be a prime ideal over p in the ring $\mathfrak{o}$ of integers. Let $\mathfrak{q}$ be a dyadic prime ideal of $\mathfrak{o}$. In convenience, if $L_{\mathfrak{p}(p)}$ represents an element n of $\mathbb{Z}_p$ then we say that $n \to L$ over $\mathbb{Z}_p$. In the dyadic case, we say that $n \to L$ over $\mathbb{Z}_2$. The following lemma will be useful for later discussion.

Lemma 1. *Let L be a primitive normal binary Hermitian lattice. Then $L_{\mathfrak{p}}$ is isometric to $\langle \epsilon, \epsilon' p^k\rangle$ for some nonnegative integer k and units ϵ, ϵ' of $\mathfrak{o}_{\mathfrak{p}}$. Moreover, $\lambda_{p^{k'}}(L_{\mathfrak{p}})$ represents all elements of $\mathbb{Z}_p$ for some nonnegative integer k'.*

Proof. Since L is primitive normal, $L_{\mathfrak{p}}$ is isometric to $\langle \epsilon, \epsilon' p^k\rangle$ for some nonnegative integer k and units ϵ, ϵ' of $\mathfrak{o}_{\mathfrak{p}}$. We may assume that any unary $\mathfrak{o}_{\mathfrak{p}}$-lattice is not isotropic. Otherwise $L_{\mathfrak{p}}$ represents all elements of $\mathbb{Z}_p$.

If $(p, m) = 1$ and $\mathfrak{p} = p\mathfrak{o}$, $k = 2\ell + r$ for some $\ell \in \mathbb{Z}$ and $r = 0, 1$, then, $\lambda_{p^{2\ell}}(L_\mathfrak{p})$ is isometric to $\langle \epsilon, \epsilon' p^r \rangle$. Since the Hermitian lattice $\langle \epsilon, \epsilon' p^r \rangle$ over $\mathfrak{o}_\mathfrak{p}$ provides the associated quadratic lattice $\langle \epsilon, \epsilon m, \epsilon' p^r, \epsilon' m p^r \rangle_{\mathbb{Z}_p} \cong \langle 1, m, p^r, m p^r \rangle_{\mathbb{Z}_p}$ over $\mathbb{Z}_p$, $\lambda_{p^{2\ell}}(L_\mathfrak{p})$ represents all elements of $\mathbb{Z}_p$.

If $(p, m) = 1$ and $\mathfrak{p} \neq p\mathfrak{o}$, then $\lambda_{p^{k-1}}(L_\mathfrak{p})$ is isometric to $\langle \epsilon, \epsilon' p \rangle$. Since the Hermitian lattice $\langle \epsilon, \epsilon' p \rangle$ provides the associated quadratic lattice $\langle \epsilon, \epsilon m, \epsilon' p, \epsilon' m p \rangle_{\mathbb{Z}_p} \cong \langle 1, m, p, mp \rangle_{\mathbb{Z}_p}$ over $\mathbb{Z}_p$, $\lambda_{p^{k-1}}(L_\mathfrak{p})$ represents all elements of $\mathbb{Z}_p$.

If $p | m$, then $\mathfrak{p} \neq p\mathfrak{o}$. So $\lambda_{p^k}(L_\mathfrak{p})$ is isometric to $\langle \epsilon, \epsilon' \rangle$. Since the Hermitian lattice $\langle \epsilon, \epsilon' \rangle$ over $\mathfrak{o}_\mathfrak{p}$ provides the associated quadratic lattice $\langle \epsilon, \epsilon m, \epsilon', \epsilon' m \rangle_{\mathbb{Z}_p} \cong \langle 1, \epsilon\epsilon', p, \epsilon\epsilon' p \rangle_{\mathbb{Z}_p}$ over $\mathbb{Z}_p$, $\lambda_{p^k}(L_\mathfrak{p})$ represents all elements of $\mathbb{Z}_p$. □

Lemma 2. *Let L be a primitive normal binary Hermitian lattice. Then $L_\mathfrak{q}$ is isometric to $\langle \epsilon, \epsilon' 2^k \rangle$ for some nonnegative integer k and units ϵ, ϵ' of $\mathfrak{o}_\mathfrak{q}$. Moreover, $\lambda_{2^{k'}}(L_\mathfrak{q})$ represents all elements of $\mathbb{Z}_p$ for some nonnegative integer k'.*

Proof. Since L is primitive normal, $L_\mathfrak{q}$ is isometric to $\langle \epsilon, \epsilon' 2^k \rangle$ for some nonnegative integer k and units ϵ, ϵ' of $\mathfrak{o}_\mathfrak{q}$.

If $m \equiv 7 \pmod 8$, then the unary Hermitian lattice $\langle \epsilon \rangle$ over $\mathfrak{o}_\mathfrak{q}$ provides the associated quadratic lattice $\begin{pmatrix} \epsilon & \epsilon/2 \\ \epsilon/2 & (m+1)\epsilon/4 \end{pmatrix}_{\mathbb{Z}_2} \cong \begin{pmatrix} 0 & 1/2 \\ 1/2 & 0 \end{pmatrix}_{\mathbb{Z}_2}$ over $\mathbb{Z}_2$. It represents all elements of $\mathbb{Z}_2$. So does $L_\mathfrak{q}$.

If $m \equiv 3 \pmod 8$, let $k = 2\ell + r$ for some $\ell \in \mathbb{Z}$ and $r = 0, 1$. Then $\lambda_{4^\ell}(L_\mathfrak{q})$ is isometric to $\langle \epsilon, \epsilon' 2^r \rangle$. Since the Hermitian lattice $\langle \epsilon, \epsilon' 2^r \rangle$ over $\mathfrak{o}_\mathfrak{q}$ provides the associated quadratic lattice $\begin{pmatrix} \epsilon & \epsilon/2 \\ \epsilon/2 & (m+1)\epsilon/4 \end{pmatrix}_{\mathbb{Z}_2} \perp \begin{pmatrix} 2^r\epsilon' & \epsilon' 2^{r-1} \\ \epsilon' 2^{r-1} & (m+1)2^{r-2}\epsilon' \end{pmatrix}_{\mathbb{Z}_2}$ which is isometric to $\begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}_{\mathbb{Z}_2} \perp \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}_{\mathbb{Z}_2}$ or $\begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}_{\mathbb{Z}_2} \perp \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}_{\mathbb{Z}_2}$ over $\mathbb{Z}_2$, $\lambda_{4^\ell}(L_\mathfrak{q})$ represents all elements of $\mathbb{Z}_2$.

If $m \equiv 1 \pmod 4$, let $k = 2\ell + r$ for some $\ell \in \mathbb{Z}$ and $r = 0, 1$. Then $\lambda_{4^\ell}(L_\mathfrak{q})$ is isometric to $\langle \epsilon, \epsilon' 2^r \rangle$. Since the Hermitian lattice $\langle \epsilon, \epsilon' 2^r \rangle$ over $\mathfrak{o}_\mathfrak{q}$ provides the associated quadratic lattice $\langle \epsilon, \epsilon m, 2^r \epsilon', 2^r \epsilon' m \rangle_{\mathbb{Z}_2}$ over $\mathbb{Z}_2$, $\lambda_{4^\ell}(L_\mathfrak{q})$ represents all elements of $\mathbb{Z}_2$.

If $m \equiv 2 \pmod 4$, then $\lambda_{2^k}(L_\mathfrak{q})$ is isometric to $\langle \epsilon, \epsilon' \rangle$. If $m = 2m'$, Hermitian lattice $\langle \epsilon, \epsilon' \rangle$ over $\mathfrak{o}_\mathfrak{q}$ provides the associated quadratic lattice $\langle \epsilon, \epsilon m, \epsilon', \epsilon' m \rangle_{\mathbb{Z}_2}$ over $\mathbb{Z}_2$. Since quadratic lattice $\langle \epsilon, \epsilon m, \epsilon', \epsilon' m \rangle_{\mathbb{Z}_2}$ is isometric to $\langle \epsilon, \epsilon', 2m'\epsilon, 2m'\epsilon' \rangle_{\mathbb{Z}_2}$, $\lambda_{2^k}(L_\mathfrak{q})$ represents all elements of $\mathbb{Z}_2$. □

Thus for all rational prime number p including 2, there is a nonnegative integer s such that $\lambda_{p^s}(L)$ represents all elements of $\mathbb{Z}_p$. So there are primes $p_1, p_2, \cdots, p_k$ and positive integers $s_1, s_2, \cdots, s_k$ such that $\widehat{L} = \lambda_{p_1^{s_1}} \circ \lambda_{p_2^{s_2}} \circ \cdots \circ \lambda_{p_k^{s_k}}(L)$ is locally universal, which means $\widehat{L}$ represents all elements of $\mathbb{Z}_p$ for all prime p. Since $\widehat{L}$ is regular, $\widehat{L}$ is universal. From the works on binary universal Hermitian lattices [7], [10], [17], we have the following proposition.

Proposition 1. *A binary normal regular lattice exists over the field $\mathbb{Q}(\sqrt{-m})$ if and only if m is*

$$1, 2, 3, 5, 6, 7, 10, 11, 15, 19, 23 \textit{ or } 31.$$

Note that the class number $h(\mathbb{Q}(\sqrt{-m}))$ is one for $m = 1, 2, 3, 7, 11, 19$ and $h(\mathbb{Q}(\sqrt{-m})) > 1$ for $m = 5, 6, 10, 15, 23, 31$.

4. CANDIDATES OF BINARY NORMAL REGULAR HERMITIAN LATTICES

In this section, we will find all candidates of binary normal regular Hermitian lattices over imaginary quadratic fields $\mathbb{Q}(\sqrt{-m})$.

If there is a rational odd prime number p_0 such that $(p_0, m) = 1$ and $L_{\mathfrak{p}_0}$ does not represent some element of $\mathbb{Z}_{p_0}$ where $\mathfrak{p}_0 = \mathfrak{p}_0(p_0)$ is a prime ideal of $\mathfrak{o}$. Let $p_1, p_2, \cdots, p_k$ be all rational prime numbers different from p_0 such that $L_{\mathfrak{p}_i}$ does not represent some element of $\mathbb{Z}_{p_i}$. Then there are positive integers $s_1, s_2, \cdots, s_k$ such that for all $i = 1, 2, \cdots, k$, $\lambda_{p_i^{s_i}}(L_{\mathfrak{p}_i})$ represents all elements of $\mathbb{Z}_{p_i}$. Let

$$\widehat{L} = \lambda_{p_1^{s_1}} \circ \lambda_{p_2^{s_2}} \circ \cdots \circ \lambda_{p_k^{s_k}}(L).$$

Then $\widehat{L}$ is regular. So $\widehat{L}_{\mathfrak{p}}$ represents all elements of $\mathbb{Z}_p$ for all prime numbers p and prime ideals $\mathfrak{p}$ except $p = p_0$. Since $\widehat{L}_{\mathfrak{p}_0}$ is primitive and normal, $\widehat{L}_{\mathfrak{p}_0}$ is isometric to $\langle \epsilon, \epsilon' p_0^k \rangle$ for some positive integer k and units ϵ, ϵ' of $\mathfrak{o}_{p_0}$. Since $(p_0, m) = 1$, $\langle \epsilon \rangle$ represents all units of $\mathbb{Z}_{p_0}$. So $\widehat{L}$ represents 1 and 2 locally. Since L is regular, $\widehat{L}$ represents 1 and 2 globally. So $\widehat{L}$ is isometric to $\langle 1 \rangle \perp M$ for some unary lattice M. If $m \neq 1, 2, 7$, then $\langle 1 \rangle$ does not represent 2. So M represents 1 or 2. Thus $\widehat{L}$ contains $\langle 1, 1 \rangle$ or $\langle 1, 2 \rangle$. Therefore $\widehat{L}$ represents all elements of $\mathbb{Z}_{p_0}$. This is a contradiction. If $m = 1$ or 7, then $\langle 1 \rangle$ cannot represent 3. Since $\widehat{L}$ is regular, 3 is not a unit of $\mathbb{Z}_{p_0}$. So $p_0 = 3$. Similarly, if $m = 2$, then $p_0 = 5$. We conclude that if $L_{\mathfrak{p}}$ does not represent some element of $\mathbb{Z}_p$, then we have following cases:

(1) $p = 2$
(2) odd prime p divides m
(3) $\begin{cases} p = 3 & \text{if } m = 1, 7, \\ p = 5 & \text{if } m = 2. \end{cases}$

To find candidates of regular lattices with efficiency, we add a condition of volume $\mathfrak{v}L$ of L as explained in the following lemma.

Lemma 3. *Let L be a binary Hermitian lattice over the imaginary quadratic field $\mathbb{Q}(\sqrt{-m})$. Let p be a rational odd prime and $\mathfrak{p} = \mathfrak{p}(p)$ be a prime ideal over p in the ring $\mathfrak{o}$ of integers and let $\mathfrak{q}$ be a dyadic prime ideal of $\mathfrak{o}$.*

(1) *If $L_{\mathfrak{p}}$ represents a unit in $\mathbb{Z}_p$ over $\mathfrak{o}_{\mathfrak{p}}$ and does not represent $p^k \epsilon$ for some nonnegative integer k, for some unit ϵ in $\mathbb{Z}_p$ over $\mathfrak{o}_{\mathfrak{p}}$, then*

$$\mathfrak{v}L \subset p^{k+1}\mathfrak{o}.$$

(2) *If $L_{\mathfrak{q}}$ represents a unit in $\mathbb{Z}_2$ over $\mathfrak{o}_{\mathfrak{q}}$ and does not represent $2^k \epsilon$ for some nonnegative integer k, for some unit ϵ in $\mathbb{Z}_2$ over $\mathfrak{o}_{\mathfrak{q}}$, then*

$$\begin{cases} \mathfrak{v}L \subset 2^{k+2}\mathfrak{o} & \textit{if } \; m \equiv 1 \pmod 4, \\ \mathfrak{v}L \subset 2^{k+3}\mathfrak{o} & \textit{if } \; m \equiv 2 \pmod 4, \\ \mathfrak{v}L \subset 2^{k+1}\mathfrak{o} & \textit{if } \; m \equiv 3 \pmod 8. \end{cases}$$

Proof. (1) Since $L_{\mathfrak{p}}$ represents a unit, $L_{\mathfrak{p}} \cong \langle a, bp^\ell \rangle$ for some units $a, b \in \mathbb{Z}_p$ and some nonnegative integer ℓ. If $\langle a \rangle$ is isotropic, then $p^k \epsilon \to \langle a \rangle$ and hence $p^k \epsilon \to L_{\mathfrak{p}}$. This is a contradiction. Therefore $\langle a \rangle$ is anisotropic. Assume that $p \nmid m$. Then the associated quadratic lattice of $L_{\mathfrak{p}}$ is isometric to $\langle a, a', bp^\ell, b'p^\ell \rangle_{\mathbb{Z}_p}$ for some units $a', b' \in \mathbb{Z}_p$. If $\ell \leq k$, then $p^k \epsilon \to L_{\mathfrak{p}}$. Now assume that $p \mid m$. Then the associated quadratic lattice is $\langle a, a'p, bp^\ell, b'p^{\ell+1} \rangle_{\mathbb{Z}}$. If $\ell \leq k$, then $p^k \epsilon \to L_{\mathfrak{p}}$. Thus $\ell \geq k+1$ and $\mathfrak{v}L_{\mathfrak{p}} = abp^\ell \mathfrak{o}_{\mathfrak{p}} \subset p^{k+1}\mathfrak{o}_{\mathfrak{p}}$.

(2) Since $L_{\mathfrak{q}}$ represents a unit in $\mathbb{Z}_2$, $L_{\mathfrak{q}}$ is isometric to $\langle a, 2^\ell b \rangle$ for some units $a, b \in \mathbb{Z}_2$ and for some integer ℓ.

Suppose $m \equiv 1 \pmod 4$. If $\ell = 0, 1$ then $L_{\mathfrak{q}} = \langle a, 2^\ell b \rangle$ represents all elements of $\mathbb{Z}_2$. Hence we have $\ell \geq 2$. Since $\lambda_{2^{\ell-1}}(L_{\mathfrak{q}}) = \langle a, 2b \rangle$ represents all elements of $\mathbb{Z}_2$, $2^{k-\ell+1}\epsilon \to \lambda_{2^{\ell-1}}(L_{\mathfrak{q}}) = \langle a, 2b \rangle$ if $\ell \leq k+1$. Hence $(2^{k-\ell+1}\epsilon)2^{\ell-1} = \epsilon 2^k \to L_{\mathfrak{q}}$, which is a contradiction. So $\ell \geq k+2$ and $\mathfrak{v}L_{\mathfrak{q}} = ab2^\ell \mathfrak{o}_{\mathfrak{q}} \subset 2^{k+2}\mathfrak{o}_{\mathfrak{q}}$.

Suppose $m \equiv 2 \pmod 4$. If $\ell = 0, 1, 2$ then $L_{\mathfrak{q}} = \langle a, 2^\ell b\rangle$ represents all elements of $\mathbb{Z}_2$. Hence we have $\ell \geq 3$. Since $\lambda_{2^{\ell-2}}(L_{\mathfrak{q}}) = \langle a, 4b\rangle$ represents all elements of $\mathbb{Z}_2$, $2^{k-\ell+2}\epsilon \to \lambda_{2^{\ell-2}}(L_{\mathfrak{q}}) = \langle a, 4b\rangle$ if $\ell \leq k+2$. Hence $(2^{k-\ell+2}\epsilon)2^{\ell-2} = \epsilon 2^k \to L_{\mathfrak{q}}$, which is a contradiction. So $\ell \geq k+3$ and $\mathfrak{v}L_{\mathfrak{q}} = ab2^\ell \mathfrak{o}_{\mathfrak{q}} \subset 2^{k+3}\mathfrak{o}_{\mathfrak{q}}$.

Suppose $m \equiv 3 \pmod 8$. If $\ell = 0, 1$, then $L_{\mathfrak{q}} = \langle a, 2^\ell b\rangle$ represents all elements of $\mathbb{Z}_2$. Hence we have $\ell \geq 2$. Since $\lambda_{2^\ell}(L_{\mathfrak{q}}) = \langle a, b\rangle$ or $\langle 2a, b\rangle$ which represents all elements of $\mathbb{Z}_2$, $2^{k-\ell}\epsilon \to \lambda_{2^\ell}(L_{\mathfrak{q}})$, if $\ell \leq k$. Hence $(2^{k-\ell}\epsilon)2^\ell = \epsilon 2^k \to L$, which is a contradiction. So $\ell \geq k+1$ and $\mathfrak{v}L_{\mathfrak{q}} = ab2^\ell\mathfrak{o} \subset 2^{k+1}\mathfrak{o}_{\mathfrak{q}}$.

$\square$

We adopt some notations from Conway-Sloane [4]. The notation p^d (resp. p^e) denotes an odd (resp. even) power of p; if $p = 2$, u_k denotes a unit of form $8n + k(k = 1, 3, 5, 7)$ and if p is odd, u_+(resp. u_-) denotes a unit which is a quadratic residue (resp. nonresidue) modulo p. Let

$$[a, \alpha, b] := \begin{pmatrix} a & \alpha \\ \overline{\alpha} & b \end{pmatrix}$$

for simplicity.

From now on, we assume L is a binary normal regular Hermitian lattice which is not universal over the imaginary quadratic field $\mathbb{Q}(\sqrt{-m})$. We begin with finding candidates of L with the information of $L_{\mathfrak{p}}$, $L_{\mathfrak{q}}$ and the volume $\mathfrak{v}L$ of L by following strategy: We assume that a is the minimum number such that $a \to \operatorname{gen} L$ and b is the minimum number such that $b \to \operatorname{gen} L$ and $b \not\to \langle a\rangle$. Then L contains a lattice $\ell = \begin{pmatrix} a & \alpha \\ \overline{\alpha} & b \end{pmatrix}$ for some $\alpha \in \mathfrak{o}$ and also $\mathfrak{v}\ell = (ab - \alpha\overline{\alpha})\mathfrak{o} \subset \mathfrak{v}L$. We call these two numbers a and b *essential numbers*. If $\mathfrak{v}\ell = \mathfrak{v}L$, then ℓ is a candidate of L. If $\mathfrak{v}\ell \subsetneq \mathfrak{v}L$, then we do more escalation which satisfies volume and rank conditions. Because most of the finding process are routine, we will give tables instead of describing in detail except special cases. In the table, all lattices are Minkowski-reduced. When we show that a binary Hermitian lattice L is not regular, we will give an integer n such that $n \to \operatorname{gen} L$ but $n \not\to L$. This number is called the *exceptional number* of L.

Case I $(m, p) = 1$.
Case I $[m = 1]$ Note that

$$u_1 \to L \iff u_5 \to L \implies 2u_1, 2u_5 \to L \text{ over } \mathbb{Z}_2;$$
$$u_3 \to L \iff u_7 \to L \implies 2u_3, 2u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \iff u_- \to L \text{ over } \mathbb{Z}_3.$$

Since L is normal, L represents a number in $\mathbb{Z}_2^\times \cap \mathbb{Z}_3^\times$. Since L is not universal, L cannot represent all elements of $\mathbb{Z}_2$ and $\mathbb{Z}_3$. According to the representability of u_1, u_3 over $\mathbb{Z}_2$ and $3u_+$ over $\mathbb{Z}_3$, we have five cases (See Table 1).
Case I $[m = 2]$ Note that

$$u_1 \to L \iff u_3 \to L \implies 2u_1, 2u_3 \to L \text{ over } \mathbb{Z}_2;$$
$$u_5 \to L \iff u_7 \to L \implies 2u_5, 2u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \iff u_- \to L \text{ over } \mathbb{Z}_5.$$

According to the representability of u_1, u_5 over $\mathbb{Z}_2$ and $5u_+$ over $\mathbb{Z}_5$, we have five cases (See Table 2).
Case I $[m = 7]$ Note that

$$u_1 \to L \iff u_3 \to L \iff u_5 \to L \iff u_7 \to L \implies 2u_1, 2u_3, 2u_5, 2u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \iff u_- \to L \text{ over } \mathbb{Z}_3.$$

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$u_1, u_3 \to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 3^2\mathfrak{o}$	$1, 7$	N.A.	
	$3u_+ \not\to L$ over $\mathbb{Z}_3$				
(2)	$u_1 \to L$, $u_3 \not\to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^2\mathfrak{o}$	$1, 21$	$\langle 1, 4\rangle$	none
	$3u_+ \to L$ over $\mathbb{Z}_3$			$\langle 1, 8\rangle$	none
				$\langle 1, 12\rangle$	6
				$\langle 1, 16\rangle$	none
				$\langle 1, 20\rangle$	6
(3)	$u_1 \to L$, $u_3 \not\to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^2{\cdot}3^2\mathfrak{o}$	$1, 77$	$\langle 1, 36\rangle$	14
	$3u_+ \not\to L$ over $\mathbb{Z}_3$			$\langle 1, 72\rangle$	28
(4-1)	$u_1 \not\to L$, $u_3, 2u_1 \to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^2\mathfrak{o}$	$2, 3$	$[2, -1+\omega, 3]$	none
	$3u_+ \to L$ over $\mathbb{Z}_3$				
(4-2)	$u_1, 2u_1 \not\to L$, $u_3 \to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$3, 7$	$[3, -1+\omega, 6]$	none
	$3u_+ \to L$ over $\mathbb{Z}_3$			$[3, 1, 3]$	none
(5-1)	$u_1 \not\to L$, $2u_1, u_3 \to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^2{\cdot}3^2\mathfrak{o}$	$2, 7$	N.A.	
	$3u_+ \not\to L$ over $\mathbb{Z}_3$				
(5-2)	$u_1, 2u_1 \not\to L$, $u_3 \to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^3{\cdot}3^2\mathfrak{o}$	$7, 11$	$[7, -2+\omega, 11]$	4
	$3u_+ \not\to L$ over $\mathbb{Z}_3$				

TABLE 1. escalation when $m = 1$

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$u_1, u_5 \to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 5^2\mathfrak{o}$	$1, 7$	N.A.	
	$5u_+ \not\to L$ over $\mathbb{Z}_5$				
(2)	$u_1 \to L$, $u_5 \not\to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^3{\cdot}5^2\mathfrak{o}$	$1, 91$	N.A.	
	$5u_+ \not\to L$ over $\mathbb{Z}_5$				
(3-1)	$u_1, 2u_5 \to L$, $u_5 \not\to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$1, 10$	$\langle 1, 8\rangle$	none
	$5u_+ \to L$ over $\mathbb{Z}_5$				
(3-2)	$u_1 \to L$, $u_5, 2u_5 \not\to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^4\mathfrak{o}$	$1, 35$	$\langle 1, 16\rangle$	none
	$5u_+ \to L$ over $\mathbb{Z}_5$			$\langle 1, 32\rangle$	none
(4)	$u_1 \not\to L$, $u_5 \to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$5, 7$	$[5, -1+\omega, 7]$	8
	$5u_+ \to L$ over $\mathbb{Z}_5$			$[5, -2+\omega, 6]$	2
				$[5, -1+2\omega, 5]$	4
				$[4, -2+4\omega, 5]$	2
				$[2, \omega, 5]$	none
(5)	$u_1 \not\to L$, $u_5 \to L$ over $\mathbb{Z}_2$	$\mathfrak{v}L \subset 2^3{\cdot}5^2\mathfrak{o}$	$7, 13$	N.A.	
	$5u_+ \not\to L$ over $\mathbb{Z}_5$				

TABLE 2. escalation when $m = 2$

So $3u_+ \not\to L$ over $\mathbb{Z}_3$. From the above conditions, $1, 5 \to L$ and $\mathfrak{v}L \subset 3^2\mathfrak{o}$. But no lattice satisfies the volume condition.

Case II $(m, p) \neq 1$.
Case II $[m = 3]$ Note that

$$u_1 \to L \Longleftrightarrow u_3 \to L \Longleftrightarrow u_5 \to L \Longleftrightarrow u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \Longrightarrow 3u_+ \to L \text{ over } \mathbb{Z}_3; \quad u_- \to L \Longrightarrow 3u_- \to L \text{ over } \mathbb{Z}_3.$$

According to the representability of $2u_1$ over $\mathbb{Z}_2$ and u_+, u_- over $\mathbb{Z}_3$, we have five cases (See Table 3).
Case II $[m = 5]$ Note that

$$u_1 \to L \Longleftrightarrow u_5 \to L \Longrightarrow 2u_3, 2u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_3 \to L \Longleftrightarrow u_7 \to L \Longrightarrow 2u_1, 2u_5 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \Longrightarrow 5u_+ \to L \text{ over } \mathbb{Z}_5; \quad u_- \to L \Longrightarrow 5u_- \to L \text{ over } \mathbb{Z}_5.$$

According to the representability of u_1, u_3 over $\mathbb{Z}_2$ and u_+, u_- over $\mathbb{Z}_5$, we have eight cases (See Table 4). Since the ring $\mathfrak{o}$ of integers is not a PID, we should also consider nonfree lattices.

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$2u_1 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 3\mathfrak{o}$	$1, 10$	$\langle 1, 3\rangle$	none
				$\langle 1, 6\rangle$	none
				$\langle 1, 9\rangle$	none
(2-1)	$2u_1 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_-, 3u_+ \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 3\mathfrak{o}$	$2, 3$	$\langle 2, 3\rangle$	none
				$[2, 1, 2]$	none
(2-2)	$2u_1 \to L$ over $\mathbb{Z}_2$ $u_+, 3u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 3^2\mathfrak{o}$	$2, 5$	$[2, 1, 5]$	none
(3)	$2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 2^2\mathfrak{o}$	$1, 10$	$\langle 1, 4\rangle$	none
(4)	$2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 2^2 \cdot 3\mathfrak{o}$	$1, 55$	$\langle 1, 12\rangle$	none
				$\langle 1, 24\rangle$	15
				$\langle 1, 36\rangle$	none
				$\langle 1, 48\rangle$	15
(5-1)	$2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_-, 3u_+ \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 2^2 \cdot 3\mathfrak{o}$	$3, 5$	$[3, 1+\omega, 5]$	none
(5-2)	$2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+, 3u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 2^2 \cdot 3^2\mathfrak{o}$	$5, 11$	$[5, 2, 8]$	none

TABLE 3. escalation when $m = 3$

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$u_1, u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 5\mathfrak{o}$	$1, 11$	$\langle 1, 5\rangle$	15
				$\langle 1, 10\rangle$	none
				$\langle 1\rangle \perp 5[2, -1+\omega, 3]$	none
(2-1)	$u_1, u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_-, 5u_+ \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 5\mathfrak{o}$	$2, 3$	$[2, 1, 3]$	11
				$[2, -1+\omega, 3] \perp \langle 5\rangle$	none
(2-2)	$u_1, u_3 \to L$ over $\mathbb{Z}_2$ $u_+, 5u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 5^2\mathfrak{o}$	$2, 3$	N.A.	
(3-1)	$u_1, 2u_1 \to L$, $u_3 \not\to \mathbb{Z}_2$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$1, 2$	N.A.	
(3-2)	$u_1 \to L$, $u_3, 2u_1 \not\to \mathbb{Z}_2$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$1, 13$	$\langle 1, 8\rangle$	none
(4-1)	$u_1, 2u_1 \to L$, $u_3 \not\to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	N.A.			
(4-2-1)	$u_1 \to L$, $u_3, 2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+, 5u_- \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3 \cdot 5\mathfrak{o}$	$1, 65$	$\langle 1, 40\rangle$	none
(4-2-2)	$u_1 \to L$, $u_3, 2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_-, 5u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3 \cdot 5^2\mathfrak{o}$	$1, 209$	$\langle 1, 200\rangle$	44
(5-1)	$u_1, 2u_1 \to L$, $u_3 \not\to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	N.A.			
(5-2-1)	$u_1 \to L$, $u_3, 2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_-, 5u_+ \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3 \cdot 5\mathfrak{o}$	$5, 13$	$\langle 5, 8\rangle$	12
(5-2-2)	$u_1 \to L$, $u_3, 2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+, 5u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3 \cdot 5^2\mathfrak{o}$	$13, 17$	$[12, 1+2\omega, 17]$	8
(6-1)	$u_1 \not\to L$, $u_3, 2u_3 \to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^2\mathfrak{o}$	$2, 3$	$[2, -1+\omega, 3] \perp \langle 4\rangle$	none
(6-2)	$u_1, 2u_3 \not\to L$, $u_3 \to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$2, 3$	$[2, -1+\omega, 3] \perp \langle 8\rangle$	8
(7-1)	$u_1 \not\to L$, $u_3, 2u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^2 \cdot 5\mathfrak{o}$	$4, 6$	$\begin{pmatrix} 4 & -2+2\omega & -2 \\ & 6 & 1+\omega \\ & & 11 \end{pmatrix}$	10
(7-2)	$u_1, 2u_3 \not\to L$, $u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	N.A.			
(8-1)	$u_1 \not\to L$, $u_3, 2u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^2 \cdot 5\mathfrak{o}$	$2, 3$	$[2, -1+\omega, 3] \perp \langle 20\rangle$	none
(8-2)	$u_1, 2u_3 \not\to L$, $u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	N.A.			

TABLE 4. escalation when $m = 5$

For the case (1), after second escalation, $L \cong \langle 1,5\rangle$ or L contains a lattice $\langle 1,10\rangle$. If L contains $\langle 1,10\rangle$, then L contains a binary lattice $\langle 1\rangle \perp \begin{pmatrix} 10 & \alpha \\ \overline{\alpha} & 5\beta \end{pmatrix}$ with $50\beta - \alpha\overline{\alpha} = 0$ and $\alpha,\, \beta \in \mathfrak{o}$. Thus $\alpha = -5+5\omega$ and $5\beta = 15$. We have candidates $\langle 1,10\rangle$ and $\langle 1\rangle \perp \begin{pmatrix} 10 & -5+5\omega \\ -5+5\overline{\omega} & 15 \end{pmatrix}$.

For the case (2-1), after second escalation, $L \cong \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}$ or L contains a unary unimodular lattice $\begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix}$ which splits L. Since $5 \to L$ and $5 \not\to \begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix}$, we can get a candidate $\begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix} \perp \langle 5\rangle$ by comparing volume of L. Similarly, we can get candidates for the cases (2-2), (6-1), (6-2) and (8-1).

For the cases (4-1), (5-1), (7-2) and (8-2), $u_+, u_- \to \lambda_{5^k}(L)$ over $\mathbb{Z}_5$ for some k by Lemma 1. Since $\lambda_{5^k}(L)$ cannot be regular by the case (3-1) or (6-2), L cannot be regular.

For the case (7-1), after second escalation, $L \cong \begin{pmatrix} 4 & 2 \\ 2 & 6 \end{pmatrix}$ or L contains a unary sublattice $2\begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix}$. The first lattice is not primitive and it is not in our consideration. In the second case, since $11 \to L$ and $11 \not\to 2\begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix}$, we conclude that $L \cong \begin{pmatrix} 4 & -2+2\omega & -2 \\ -2+2\overline{\omega} & 6 & 1+\omega \\ -2 & 1+\overline{\omega} & 11 \end{pmatrix}$ with $\mathfrak{v}L = 20\mathfrak{o}$.

Case II $[m=6]$ Note that

$$u_1 \to L \Longleftrightarrow u_7 \to L \Longrightarrow 2u_3, 2u_5 \to L \text{ over } \mathbb{Z}_2;$$
$$u_3 \to L \Longleftrightarrow u_5 \to L \Longrightarrow 2u_1, 2u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \Longrightarrow 3u_- \to L \text{ over } \mathbb{Z}_3; \quad u_- \to L \Longrightarrow 3u_+ \to L \text{ over } \mathbb{Z}_3.$$

According to the representability of u_1, u_3 over $\mathbb{Z}_2$ and u_+, u_- over $\mathbb{Z}_3$, we have eight cases (See Table 5). Since the ring $\mathfrak{o}$ of integers is not a PID, we should also consider nonfree lattices.

For the case (2-2), after second escalation, $L \cong \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$ or L contains $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$ by the volume condition. For the second case, note that $6 \to L$ and $6 \not\to \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$. Since $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$ is a unary sublattice which splits L, $L \cong \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix} \perp \langle 3\rangle$ or L contains a lattice $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix} \perp \langle 6\rangle$ by the volume condition. Since 9 is an exceptional number of $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix} \perp \langle 3\rangle$, it is not regular. For the last case note that $15 \to L$ and $15 \not\to \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix} \perp \langle 6\rangle$. So L contains $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix} \perp \begin{pmatrix} 6 & \delta \\ \overline{\delta} & 3\rho \end{pmatrix}$ with $18\rho - \delta\overline{\delta} = 0$ with $\delta, \rho \in \mathfrak{o}$. Then we have a candidate $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix} \perp 3\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$ with volume $3\mathfrak{o}$. Note that this lattice is isometric to the *free* lattice $\begin{pmatrix} 9 & 4\omega \\ 4\overline{\omega} & 11 \end{pmatrix}$. Similarly, we can get candidates for the cases (2-1) and (6).

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1-1)	$u_1, u_3 \to L$ over $\mathbb{Z}_2$ $u_+, 3u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 3\mathfrak{o}$	$1, 3$	$\langle 1, 3\rangle$	none
(1-2)	$u_1, u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_-, 3u_+ \not\to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 3^2\mathfrak{o}$	$1, 13$	$\langle 1, 9\rangle$	27
(2-1)	$u_1, u_3 \to L$ over $\mathbb{Z}_2$ $u_+, 3u_- \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 3^2\mathfrak{o}$	$2, 3$	$[2, \omega, 3] \perp \langle 9\rangle$	26
(2-2)	$u_1, u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_-, 3u_- \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 3\mathfrak{o}$	$2, 3$	$[2, 0, 3]$ $[2, \omega, 3] \perp \langle 3\rangle$ $[2, \omega, 3] \perp 3[2, \omega, 3]$	6 9 none
(3-1)	$u_1, 2u_1 \to L$, $u_3 \not\to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$1, 2$	N.A.	
(3-2)	$u_1 \to L$, $2u_1, u_3 \not\to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 2^4\mathfrak{o}$	$1, 17$	$\langle 1, 16\rangle$	7
(4)	$u_1 \to L$, $u_3 \not\to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$	N.A.			
(5)	$u_1 \to L$, $u_3 \not\to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$	N.A.			
(6)	$u_1 \not\to L$, $u_3 \to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_3$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$2, 3$	$[2, \omega, 3] \perp \langle 8\rangle$	6
(7)	$u_1 \not\to L$, $u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$	N.A.			
(8)	$u_1 \not\to L$, $u_3 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$	N.A.			

TABLE 5. escalation when $m = 6$

For the case (4), (5), (7) and (8), $u_+, u_- \to \lambda_{3^k}(L)$ over $\mathbb{Z}_3$ for some k by Lemma 1. since $\lambda_{3^k}(L)$ cannot be regular by the case (3-1), (3-2) or (6), L cannot be regular.

Case II $[m = 7]$ Note that

$$u_1 \to L \iff u_3 \to L \iff u_5 \to L \iff u_7 \to L \implies 2u_1, 2u_3, 2u_5, 2u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \implies 7u_+ \to L \text{ over } \mathbb{Z}_7; \quad u_- \to L \implies 7u_- \to L \text{ over } \mathbb{Z}_7.$$

According to the representability of u_+, u_- over $\mathbb{Z}_7$, we have two cases (See Table 6).

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_7$	$\mathfrak{v}L \subset 7\mathfrak{o}$	$1, 15$	$\langle 1, 7\rangle$ $\langle 1, 14\rangle$	none none
(2)	$u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_7$	$\mathfrak{v}L \subset 7\mathfrak{o}$	$3, 5$	$[3, 1, 5]$ $[3, \omega, 3]$	7 none

TABLE 6. escalation when $m = 7$

Case II $[m = 10]$ Note that

$$u_1 \to L \iff u_3 \to L \implies 2u_5, 2u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_5 \to L \iff u_7 \to L \implies 2u_1, 2u_3 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \implies 5u_- \to L \text{ over } \mathbb{Z}_5; \quad u_- \to L \implies 5u_+ \to L \text{ over } \mathbb{Z}_5.$$

According to the representability of u_1, u_5 over $\mathbb{Z}_2$ and u_+, u_- over $\mathbb{Z}_7$, we have eight cases (See Table 7). Since the ring $\mathfrak{o}$ of integers is not a PID, we should also consider nonfree lattices.

For the case (6), L contains a unimodular sublattice $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 5 \end{pmatrix}$ which splits L. Since $6 \to L$ and $6 \not\to \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 5 \end{pmatrix}$, L contains $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 5 \end{pmatrix} \perp \langle a\rangle$ with $a \leq 6$. But this lattice cannot have the volume contained in $8\mathfrak{o}$. So we have no candidates.

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$u_1, u_5 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 5\mathfrak{o}$	$1, 6$	$\langle 1, 5\rangle$	none
(2)	$u_1, u_5 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 5\mathfrak{o}$	$2, 3$	$[2, 1, 3]$	5
(3)	$u_1 \to L$, $u_5 \not\to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$1, 3$	N.A.	
(4)	$u_1 \to L$, $u_5 \not\to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	N.A.			
(5)	$u_1 \to L$, $u_5 \not\to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	N.A.			
(6)	$u_1 \not\to L$, $u_5 \to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 2^3\mathfrak{o}$	$2, 5$	N.A.	
(7)	$u_1 \not\to L$, $u_5 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	N.A.			
(8)	$u_1 \not\to L$, $u_5 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	N.A.			

Table 7. escalation when $m = 10$

For the case (4), (5), (7) and (8), $u_+, u_- \to \lambda_{5^k}(L)$ over $\mathbb{Z}_3$ for some k by Lemma 1. Since $\lambda_{5^k}(L)$ cannot be regular by the case (3) or (6), L cannot be regular.

Case II $[m = 11]$ Note that

$$u_1 \to L \Longleftrightarrow u_3 \to L \Longleftrightarrow u_5 \to L \Longleftrightarrow u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \Longrightarrow 11u_+ \to L \text{ over } \mathbb{Z}_{11}; \quad u_- \to L \Longrightarrow 11u_- \to L \text{ over } \mathbb{Z}_{11}.$$

According to the representability of $2u_1$ over $\mathbb{Z}_2$ and u_+, u_- over $\mathbb{Z}_7$, we have five cases (See Table 8).

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$2u_1 \to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_{11}$	$\mathfrak{v}L \subset 11\mathfrak{o}$	$1, 14$	$\langle 1, 11\rangle$	none
(2)	$2u_1 \to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_{11}$	$\mathfrak{v}L \subset 11\mathfrak{o}$	$2, 7$	$[2, \omega, 7]$	11
(3)	$2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+, u_- \to L$ over $\mathbb{Z}_{11}$	$\mathfrak{v}L \subset 2^2\mathfrak{o}$	$1, 7$	$\langle 1, 4\rangle$	none
(4)	$2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_{11}$	$\mathfrak{v}L \subset 2^2{\cdot}11\mathfrak{o}$	$1, 91$	$\langle 1, 44\rangle$ $\langle 1, 88\rangle$	none 77
(5)	$2u_1 \not\to L$ over $\mathbb{Z}_2$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_{11}$	$\mathfrak{v}L \subset 2^2{\cdot}11\mathfrak{o}$	$7, 13$	$[7, \omega, 13]$ $[7, 2\omega, 8]$	8 11

Table 8. escalation when $m = 11$

Case II $[m = 15]$ Note that

$$u_1 \to L \Longleftrightarrow u_3 \to L \Longleftrightarrow u_5 \to L \Longleftrightarrow u_7 \to L \Longrightarrow 2u_1, 2u_3, 2u_5, 2u_7 \to L \text{ over } \mathbb{Z}_2;$$
$$u_+ \to L \Longrightarrow 3u_- \to L \text{ over } \mathbb{Z}_3; \quad u_- \to L \Longrightarrow 3u_+ \to L \text{ over } \mathbb{Z}_3;$$
$$u_+ \to L \Longrightarrow 5u_- \to L \text{ over } \mathbb{Z}_5; \quad u_- \to L \Longrightarrow 5u_+ \to L \text{ over } \mathbb{Z}_5.$$

According to the representability of u_+, u_- over $\mathbb{Z}_5$ and u_+, u_- over $\mathbb{Z}_7$, we have eight cases (See Table 9).

For the case (1), $L \cong \langle 1, 5\rangle$ or L contains $\langle 1, 10\rangle$. Since 5 is an exceptional number of $\langle 1, 10\rangle$, L contains a lattice $\langle 1\rangle \perp \begin{pmatrix} 10 & \alpha \\ \overline{\alpha} & 5\beta \end{pmatrix}$ with $50\beta - \alpha\overline{\alpha} = 0$ and α, $\beta \in \mathfrak{o}$. Thus $L \cong \langle 1\rangle \perp 5\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$.

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$u_+, u_- \to L$ over $\mathbb{Z}_3$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 5\mathfrak{o}$	1, 11	$\langle 1, 5\rangle$ $\langle 1, 10\rangle$ $\langle 1\rangle \perp 5[2, \omega, 5]$	none 5 5
(2)	$u_+, u_- \to L$ over $\mathbb{Z}_3$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 5\mathfrak{o}$	2, 3	$[2, 1, 3]$ $[2, \omega, 2] \perp \langle 5\rangle$	5 none
(3)	$u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3\mathfrak{o}$	1, 7	$\langle 1, 3\rangle$ $\langle 1, 6\rangle$	none 3
(4-1)	$u_+, 3u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3 \cdot 5\mathfrak{o}$	1, 21	$\langle 1, 15\rangle$	45
(4-2)	$u_+ \to L$, $u_-, 3u_+ \not\to L$ over $\mathbb{Z}_3$ $u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3^2 \cdot 5\mathfrak{o}$	1, 91	$\langle 1, 45\rangle$ $\langle 1, 90\rangle$	17 145
(5-1-1)	$u_+, 3u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$ $u_+ \not\to L$, $u_-, 5u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3 \cdot 5\mathfrak{o}$	3, 7	$[3, 1+\omega, 7]$	15
(5-1-2)	$u_+, 3u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_3$ $u_+, 5u_- \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3 \cdot 5^2\mathfrak{o}$	3, 7	N.A.	
(5-2-1)	$u_+ \to L$, $u_-, 3u_+ \not\to L$ over $\mathbb{Z}_3$ $u_+ \not\to L$, $u_-, 5u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3^2 \cdot 5\mathfrak{o}$	7, 11	$[7, 2, 7]$	13
(5-2-2)	$u_+ \to L$, $u_-, 3u_+ \not\to L$ over $\mathbb{Z}_3$ $u_+, 5u_- \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3^2{\cdot}5^2\mathfrak{o}$	7, 13	N.A.	
(6-1)	$u_+ \not\to L$, $u_-, 3u_- \to L$ over $\mathbb{Z}_3$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3\mathfrak{o}$	2, 6	$[2, \omega, 2] \perp \langle 6\rangle$ $[2, \omega, 2] \perp 3[2, \omega, 2]$ $\cong [8, -1+4\omega, 8]$	15 none
(6-2)	$u_+, 3u_- \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$ $u_+, u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3^2\mathfrak{o}$	2, 3	$[2, \omega, 2] \perp \langle 9\rangle$	none
(7-1-1)	$u_+ \not\to L$, $u_-, 3u_- \to L$ over $\mathbb{Z}_3$ $u_+, 5u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3 \cdot 5\mathfrak{o}$	5, 6	$[5, -1+2\omega, 6]$ $\langle 5\rangle \perp 3[2, \omega, 2]$	9 21
(7-1-2)	$u_+ \not\to L$, $u_-, 3u_- \to L$ over $\mathbb{Z}_3$ $u_+ \to L$, $u_-, 5u_+ \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3 \cdot 5^2\mathfrak{o}$	6, 11	N.A.	
(7-2-1)	$u_+, 3u_- \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$ $u_+, 5u_+ \to L$, $u_- \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3^2 \cdot 5\mathfrak{o}$	5, 11	$[5, 2+\omega, 11]$	9
(7-2-2)	$u_+, 3u_- \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$ $u_+ \to L$, $u_-, 5u_+ \not\to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3^2{\cdot}5^2\mathfrak{o}$	11, 14	N.A.	
(8-1)	$u_+ \not\to L$, $u_-, 3u_- \to L$ over $\mathbb{Z}_3$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3 \cdot 5\mathfrak{o}$	2, 3	N.A.	
(8-2)	$u_+, 3u_- \not\to L$, $u_- \to L$ over $\mathbb{Z}_3$ $u_+ \not\to L$, $u_- \to L$ over $\mathbb{Z}_5$	$\mathfrak{v}L \subset 3^2 \cdot 5\mathfrak{o}$	2, 3	$[2, \omega, 2] \perp \langle 45\rangle$	35

TABLE 9. escalation when $m = 15$

For the case (2), after second escalation, $L \cong \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}$ or L contains a unary unimodular lattice $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$ which splits L. Since $7 \to L$ and $7 \not\to \begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix}$, we can get candidate $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix} \perp \langle 5\rangle$ by comparing volume of L. Similarly, we can get results for the cases (6-2), (8-1) and (8-2).

For the case (3), $L \cong \langle 1, 3\rangle$ or L contains $\langle 1, 6\rangle$. When L contains $\langle 1, 6\rangle$, since $3 \to \operatorname{gen}\langle 1, 6\rangle$, $3 \to L$ and hence we have a candidate $\langle 1, 3\rangle$ by the volume condition. Similarly, for the case (4-2), L contains $\langle 1, 90\rangle$ or $L \cong \langle 1, 45\rangle$. If L contains $\langle 1, 90\rangle$, then since $45 \to \operatorname{gen}\langle 1, 90\rangle$ and $45 \not\to \langle 1, 90\rangle$, $L \cong \langle 1, 45\rangle$ by the volume condition.

For the case (6-1), after second escalation, L contains $\langle 2, 6\rangle$, $\begin{pmatrix} 2 & 1+\omega \\ 1+\overline{\omega} & 3 \end{pmatrix}$ or $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$. If L contains $\langle 2, 6\rangle$, from the condition $3 \to L$, L contains a lattice $\begin{pmatrix} 2 & 0 & \beta \\ 0 & 6 & \gamma \\ \overline{\beta} & \overline{\gamma} & 3 \end{pmatrix}$ with $3 \mid (6 - \beta\overline{\beta})$, $3 \mid (18 - \gamma\overline{\gamma})$ and its determinant $36 - 6\beta\overline{\beta} - 2\gamma\overline{\gamma} = 0$.

So L contains a lattice $\begin{pmatrix} 2 & 1+\omega \\ 1+\overline{\omega} & 3 \end{pmatrix} \perp \langle 6 \rangle \cong \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix} \perp \langle 6 \rangle$. Since $15 \rightarrow L$ and $15 \not\rightarrow \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix} \perp \langle 6 \rangle$, we have a candidate $L \cong \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix} \perp 3\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$, via similar way. Note that L is isometric to the binary *free* lattice $\begin{pmatrix} 8 & -3+4\omega \\ -3+4\overline{\omega} & 8 \end{pmatrix}$. If L contains $\begin{pmatrix} 2 & 1+\omega \\ 1+\overline{\omega} & 3 \end{pmatrix}$ or $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, then we know that there is no candidates via similar way.

Case II $[m = 19]$ Note that

$$u_1 \rightarrow L \Longleftrightarrow u_3 \rightarrow L \Longleftrightarrow u_5 \rightarrow L \Longleftrightarrow u_7 \rightarrow L \text{ over } \mathbb{Z}_2;$$
$$u_+ \rightarrow L \Longrightarrow 19u_+ \rightarrow L \text{ over } \mathbb{Z}_{19}; \quad u_- \rightarrow L \Longrightarrow 19u_- \rightarrow L \text{ over } \mathbb{Z}_{19}.$$

According to the representability of $2u_1$ over $\mathbb{Z}_2$ and u_+, u_- over $\mathbb{Z}_{19}$, we have two cases (See Table 10).

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1-1)	$2u_1 \rightarrow L$ over $\mathbb{Z}_2$ $u_+ \rightarrow L$, $u_- \not\rightarrow L$ over $\mathbb{Z}_{19}$	$\mathfrak{v}L \subset 19\mathfrak{o}$	1, 6	N.A.	
(1-2)	$2u_1 \not\rightarrow L$ over $\mathbb{Z}_2$ $u_+ \rightarrow L$, $u_- \not\rightarrow L$ over $\mathbb{Z}_{19}$	$\mathfrak{v}L \subset 2^2{\cdot}19\mathfrak{o}$	1, 39	N.A.	
(2-1)	$2u_1 \rightarrow L$ over $\mathbb{Z}_2$ $u_+ \not\rightarrow L$, $u_- \rightarrow L$ over $\mathbb{Z}_{19}$	$\mathfrak{v}L \subset 19\mathfrak{o}$	2, 3	N.A.	
(2-2)	$2u_1 \not\rightarrow L$ over $\mathbb{Z}_2$ $u_+ \not\rightarrow L$, $u_- \rightarrow L$ over $\mathbb{Z}_{19}$	$\mathfrak{v}L \subset 2^2{\cdot}19\mathfrak{o}$	3, 13	N.A.	

TABLE 10. escalation when $m = 19$

Case II $[m = 23]$ Note that

$$u_1 \rightarrow L \Longleftrightarrow u_3 \rightarrow L \Longleftrightarrow u_5 \rightarrow L \Longleftrightarrow u_7 \rightarrow L \Longrightarrow 2u_1, 2u_3, 2u_5, 2u_7 \rightarrow L \text{ over } \mathbb{Z}_2;$$
$$u_+ \rightarrow L \Longrightarrow 23u_+ \rightarrow L \text{ over } \mathbb{Z}_{23}; \quad u_- \rightarrow L \Longrightarrow 23u_- \rightarrow L \text{ over } \mathbb{Z}_{23}.$$

According to the representability of u_+, u_- over $\mathbb{Z}_{23}$, we have two cases (See Table 11).

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$u_+ \rightarrow L$, $u_- \not\rightarrow L$ over $\mathbb{Z}_{23}$	$\mathfrak{v}L \subset 23\mathfrak{o}$	1, 2	N.A.	
(2)	$u_+ \not\rightarrow L$, $u_- \rightarrow L$ over $\mathbb{Z}_{23}$	$\mathfrak{v}L \subset 23\mathfrak{o}$	5, 7	$[5, 2+\omega, 7]$	10

TABLE 11. escalation when $m = 23$

Case II $[m = 31]$ Note that

$$u_1 \rightarrow L \Longleftrightarrow u_3 \rightarrow L \Longleftrightarrow u_5 \rightarrow L \Longleftrightarrow u_7 \rightarrow L \Longrightarrow 2u_1, 2u_3, 2u_5, 2u_7 \rightarrow L \text{ over } \mathbb{Z}_2;$$
$$u_+ \rightarrow L \Longrightarrow 31u_+ \rightarrow L \text{ over } \mathbb{Z}_{31}; \quad u_- \rightarrow L \Longrightarrow 31u_- \rightarrow L \text{ over } \mathbb{Z}_{31}.$$

According to the representability of u_+, u_- over $\mathbb{Z}_{31}$, we have two cases (See Table 12).

Local Condition		Volume	Ess.#	Reduced Lattice	Exc.#
(1)	$u_+ \rightarrow L$, $u_- \not\rightarrow L$ over $\mathbb{Z}_{31}$	$\mathfrak{v}L \subset 31\mathfrak{o}$	1, 2	N.A.	
(2)	$u_+ \not\rightarrow L$, $u_- \rightarrow L$ over $\mathbb{Z}_{31}$	$\mathfrak{v}L \subset 31\mathfrak{o}$	3, 6	N.A.	

TABLE 12. escalation when $m = 31$

5. Complete list of binary regular Hermitian lattices

In this section, we will prove the regularity of each candidate. If its class number is one, then it is trivially regular. We know that the class numbers of following lattices are one and hence they are regular. Some diagonal lattices are checked by [20], and the other nondiagonal lattices are checked by authors via comparing discriminants and local properties.

field	class number one lattices
$\mathbb{Q}(\sqrt{-1})$	$\langle 1,4\rangle$, $\begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix}$, $\begin{pmatrix} 3 & -1+\omega \\ -1+\overline{\omega} & 6 \end{pmatrix}$, $\begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$
$\mathbb{Q}(\sqrt{-3})$	$\langle 1,3\rangle$, $\langle 1,4\rangle$, $\langle 1,6\rangle$, $\langle 2,3\rangle$, $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, $\begin{pmatrix} 2 & 1 \\ 1 & 5 \end{pmatrix}$, $\begin{pmatrix} 3 & 1+\omega \\ 1+\overline{\omega} & 5 \end{pmatrix}$, $\begin{pmatrix} 5 & 2 \\ 2 & 8 \end{pmatrix}$
$\mathbb{Q}(\sqrt{-7})$	$\begin{pmatrix} 3 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$

Now, we prove all the surviving candidates are actually regular. Most of proofs are using a ternary regular sublattice whose class number one. Whereas in the the proof of Case II $[m=7]$ (2), we give an efficient bound for the numbers represented by a specific quaternary quadratic form. This new arithmetic method uses ternary quadratic forms which are not regular.

Case I $[m = 1]$

(1) $L = \langle 1, 8\rangle$ is regular over $\mathbb{Q}(\sqrt{-1})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 1 \pmod 4 \text{ or } n \equiv 2 \pmod 8\}.$$

Since $\langle 2, 8\rangle = 2\langle 1, 4\rangle$ is a sublattice of L, if $n \to \langle 1, 4\rangle$, then $2n \to \langle 2, 8\rangle$ and hence $2n \to L$. From the Case I $[m = 1]$ (1), we know that $\langle 1, 4\rangle$ is regular, which represents all positive integers n such that $n \equiv 0, 1, 2 \pmod 4$. Hence L represents all positive integers n such that $n \equiv 0 \pmod 4$ or $n \equiv 2 \pmod 8$. On the other hand, the associated quadratic lattice of L is

$$x\overline{x} + 8y\overline{y} = x_1^2 + x_2^2 + 8y_1^2 + 8y_2^2.$$

Since it has a regular sublattice $x_1^2 + x_2^2 + 8y_1^2$ [12], which represents all positive integers $n \equiv 1 \pmod 4$, L represents all positive integers $n \equiv 1 \pmod 4$. Therefore L is regular. □

(2) $L = \langle 1, 16\rangle$ is regular over $\mathbb{Q}(\sqrt{-1})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 1 \pmod 4, n \equiv 0, 2 \pmod 8 \text{ or } n \equiv 4 \pmod{16}\}.$$

Since $\langle 2, 16\rangle = 2\langle 1, 8\rangle$ is a sublattice of L, if $n \to \langle 1, 8\rangle$, then $2n \to \langle 2, 16\rangle$ and hence $2n \to L$. From the Case I $[m = 1]$ (2), we know that $\langle 1, 8\rangle$ is regular, which represents all positive integers n such that $n \equiv 0, 1 \pmod 4$ or $n \equiv 2 \pmod 8$. Hence L represents all positive integers n such that $n \equiv 0, 2 \pmod 4$ or $n \equiv 4 \pmod{16}$. On the other hand, the associated quadratic lattice of L is

$$x\overline{x} + 16y\overline{y} = x_1^2 + x_2^2 + 16y_1^2 + 16y_2^2.$$

Since it has a regular sublattice $x_1^2 + x_2^2 + 16y_1^2$ [12], which represents all positive integers $n \equiv 1 \pmod 4$, L represents all positive integers $n \equiv 1 \pmod 4$. Therefore L is regular. □

Case I $[m = 2]$

(1) $L = \langle 1, 8\rangle$ is regular over $\mathbb{Q}(\sqrt{-2})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 1, 3 \pmod 8 \text{ or } n \equiv 0 \pmod 2\}.$$

Since $\langle 2, 8\rangle = 2\langle 1, 4\rangle$ is a sublattice of L and $\langle 1, 4\rangle$ is universal [7], $n \to L$ for all $n \equiv 0 \pmod 2$. On the other hand, the associated quadratic form of L is

$$x\overline{x} + 8y\overline{y} = x_1^2 + 2x_2^2 + 8y_1^2 + 16y_2^2.$$

Since it has a regular sublattice $x_1^2 + 2x_2^2 + 8y_1^2$ [12], which represents all positive integers $n \equiv 1, 3 \pmod 8$, these n are all represented by L. Therefore L is regular. □

(2) $L = \langle 1, 16\rangle$ is regular over $\mathbb{Q}(\sqrt{-2})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0 \pmod 4, n \equiv 1, 3 \pmod 8 \text{ or } n \equiv 2, 6 \pmod{16}\}.$$

Since $\langle 4, 16\rangle = 4\langle 1, 4\rangle$ is a sublattice of L and $\langle 1, 4\rangle$ is universal [7], $n \to L$ for all $n \equiv 0 \pmod 4$. On the other hand, the associated quadratic form of L is

$$x\overline{x} + 8y\overline{y} = x_1^2 + 2x_2^2 + 16y_1^2 + 32y_2^2.$$

Since it has a regular sublattice $x_1^2 + 2x_2^2 + 16y_1^2$ [12], which represents all positive integers $n \equiv 1, 3 \pmod 8$ or $n \equiv 2, 6 \pmod{16}$, these n are all represented by L. Therefore L is regular. □

(3) $L = \langle 1, 32\rangle$ is regular over $\mathbb{Q}(\sqrt{-2})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 1, 3 \pmod 8, n \equiv 2, 6 \pmod{16} \text{ or } n \equiv 4, 12 \pmod{32}\}.$$

Since $\langle 8, 32\rangle = 8\langle 1, 4\rangle$ is a sublattice of L and $\langle 1, 4\rangle$ is universal [7], $n \to L$ for all $n \equiv 0 \pmod 8$. On the other hand, the associated quadratic form of L is

$$x\overline{x} + 8y\overline{y} = x_1^2 + 2x_2^2 + 32y_1^2 + 64y_2^2.$$

Since it has a regular sublattice $x_1^2 + 2x_2^2 + 32y_1^2$ [12], which represents all positive integers $n \equiv 1, 3 \pmod 8$, $n \equiv 2, 6 \pmod{16}$ or $n \equiv 4, 12 \pmod{32}$, these n are all represented by L. Therefore L is regular. □

(4) $L = \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 5 \end{pmatrix}$ is regular over $\mathbb{Q}(\sqrt{-2})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0 \pmod 2, \text{ or } n \equiv 5, 7 \pmod 8\}.$$

Consider a universal lattice $\langle 1, 4\rangle$ with basis $\{v_1, v_2\}$ [7]. We know that L is a sublattice $\langle 1, 4\rangle$ with basis $\{\omega v_1, v_1 + v_2\}$. If $n \equiv 0 \pmod 2$, then $n \to L$, since L contains a sublattice $\langle 2, 8\rangle = 2\langle 1, 4\rangle$ and $\langle 1, 4\rangle$ is universal. If $n \equiv 5, 7 \pmod 8$, then $n \to \langle 1, 4\rangle$ by universality of $\langle 1, 4\rangle$ and hence

$$n = H(xv_1 + yv_2) = x\overline{x} + 4y\overline{y} = x_1^2 + 2x_2^2 + 4y_1^2 + 8y_2^2$$

has a solution for some $x = x_1 + \omega x_2$, $y = y_1 + \omega y_2 \in \mathfrak{o}$. Since $n \equiv 5, 7 \pmod 8$, $x_1 \equiv y_1 \equiv 1 \pmod 2$. So $x_1 - y_1$ is even and multiple of ω. We deduce that

$$xv_1 + yv_2 = (x_1 - y_1)v_1 + \omega(x_2 - y_2)v_1 + (y_1 + \omega y_2)(v_1 + v_2) \in L$$

and hence $n \to L$. Therefore L is regular. □

Case II $[m = 3]$

(1) $L = \langle 1, 9\rangle$ is regular over $\mathbb{Q}(\sqrt{-3})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 1 \pmod 3 \text{ or } n \equiv 0, 3 \pmod 9\}.$$

Since $\langle 3, 9\rangle = 3\langle 1, 3\rangle$ is a sublattice of L, if $n \to \langle 1, 3\rangle$, then $3n \to \langle 3, 9\rangle$ and hence $3n \to L$. From Case II [$m = 3$] (1), we know that $\langle 1, 3\rangle$ is regular, which represents all positive integers n such that $n \equiv 0, 1 \pmod 3$. Hence L represents all positive integers n such that $n \equiv 0, 3 \pmod 9$. On the other hand, the associated quadratic lattice of L is

$$x\overline{x} + 9y\overline{y} = x_1^2 + x_1x_2 + x_2^2 + 9y_1^2 + 9y_1y_2 + 9y_2^2.$$

Since it has a regular sublattice $x_1^2 + x_1x_2 + x_2^2 + 9y_1^2$ [12], which represents all positive integers $n \equiv 1 \pmod 3$, these n are all represented by L. Therefore L is regular. □

(2) $L = \langle 1, 12\rangle$ is regular over $\mathbb{Q}(\sqrt{-3})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 1, 3, 4, 7, 9 \pmod{12}\}.$$

Consider regular lattices $\langle 1, 3\rangle$ and $\langle 1, 4\rangle$ (see Case II [$m = 3$] (1), (2)). Since $\langle 1, 3\rangle$ represents all positive integers n such that $n \equiv 0, 1 \pmod 3$ and L contains a sublattice $\langle 4, 12\rangle = 4\langle 1, 3\rangle$, L represents all positive integers n such that $n \equiv 0, 4 \pmod{12}$. Since $\langle 1, 4\rangle$ represents all positive integers n such that $n \equiv 0, 1, 3 \pmod 4$ and L contains a sublattice $\langle 3, 12\rangle = 3\langle 1, 4\rangle$, L represents all positive integers n such that $n \equiv 0, 1, 7 \pmod{12}$. On the other hand, the associated quadratic lattice of L is

$$x\overline{x} + 12y\overline{y} = x_1^2 + x_1x_2 + x_2^2 + 12y_1^2 + 12y_1y_2 + 12y_2^2.$$

Since it contains a regular lattice $x_1^2 + x_1x_2 + x_2^2 + 12y_1^2$ [12], which represents all positive integers $n \equiv 0, 1 \pmod 3$, these n are all represented by L. Hence L is regular. □

(3) $L = \langle 1, 36\rangle$ is regular over $\mathbb{Q}(\sqrt{-3})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{Z} \mid n \equiv 0, 1, 3 \pmod 4 \text{ and } n \equiv 0, 1 \pmod 3\}.$$

Consider regular lattices $\langle 1, 9\rangle$, $\langle 1, 12\rangle$ (see Case II [$m = 3$] (5), (6)). Since L contains sublattices $\langle 4, 36\rangle = 4\langle 1, 9\rangle$ and $\langle 3, 36\rangle = 3\langle 1, 12\rangle$, L represents all positive integers n such that $n \equiv 0 \pmod 4$ and $n \equiv 0, 1 \pmod 3$, or $n \equiv 0 \pmod 3$ and $n \equiv 1 \pmod 2$. So it suffices to show that L represents all positive integers n such that n is odd and $n \equiv 1 \pmod 3$. The associated quadratic lattice of L is

$$x\overline{x} + 36y\overline{y} = x_1^2 + x_1x_2 + x_2^2 + 36y_1^2 + 36y_1y_2 + 36y_2^2,$$

and it contains a sublattice isometric to $\langle 1, 3, 36, 108\rangle_{\mathbb{Z}} = \langle 1\rangle_{\mathbb{Z}} \perp 3\langle 1, 12, 36\rangle_{\mathbb{Z}}$. Since $\langle 1, 12, 36\rangle_{\mathbb{Z}}$ is regular [12], $3\langle 1, 12, 36\rangle_{\mathbb{Z}} = \langle 3, 36, 108\rangle_{\mathbb{Z}}$ represents all positive integer $n \equiv 3, 12 \pmod{36}$. If $n \equiv 1 \pmod{12}$ and $n \geq 49$, then $n - a^2 \equiv 12 \pmod{36}$ for $a = 1, 5, 7$ and hence L represents n. If $n \equiv 7 \pmod{12}$ and $n \geq 64$, then $n - a^2 \equiv 3 \pmod{36}$ for $a = 2, 4, 8$ and hence L represents n. It is an easy work to check $n \to L$ for $1, 7, 13, 19, 25, 31, 37, 43, 49, 55$. Therefore L is regular. □

Case II [$m = 5$]

(1) $L = \langle 1, 8\rangle$ is regular over $\mathbb{Q}(\sqrt{-5})$.

Proof. Note that

$$H(\operatorname{gen} L)=\{n\in\mathbb{N}_0 \mid n\equiv 0,1 \pmod 4 \text{ or } n\equiv 6 \pmod 8\}.$$

Since $\langle 4,8\rangle = 4\langle 1,2\rangle$ is a sublattice of L and $\langle 1,2\rangle$ is universal [10], $n\to L$ for all $n\equiv 0 \pmod 4$. On the other hand, the associated quadratic lattice of L is

$$x\overline{x}+8y\overline{y}=x_1^2+5x_2^2+8y_1^2+40y_2^2.$$

Since it has a regular sublattice $x_1^2+5x_2^2+8y_1^2$ [12], which represents all positive integers $n\equiv 1 \pmod 4$ or $n\equiv 6 \pmod 8$, these all n are represented by L. Therefore L is regular. □

(2) $L=\langle 1,10\rangle$ is regular over $\mathbb{Q}(\sqrt{-5})$.

Proof. Note that

$$H(\operatorname{gen} L)=\{n\in\mathbb{N}_0 \mid n\equiv 0,1,4 \pmod 5\}.$$

Since $\langle 5,10\rangle = 5\langle 1,2\rangle$ is a sublattice of L and $\langle 1,2\rangle$ is universal [10], $n\to L$ for all $n\equiv 0 \pmod 5$. On the other hand, the associated quadratic lattice of L is

$$x\overline{x}+10y\overline{y}=x_1^2+5x_2^2+10y_1^2+50y_2^2.$$

Since it has a regular sublattice $x_1^2+5x_2^2+10y_1^2$ [12], which represents all positive integers $n\equiv 1,4 \pmod 5$, $n\to L$ for all positive integers $n\equiv 1,4 \pmod 5$. Therefore L is regular. □

(3) $L=\langle 1\rangle\perp 5\begin{pmatrix}2 & -1+\omega\\ -1+\overline{\omega} & 3\end{pmatrix}$ is regular over $\mathbb{Q}(\sqrt{-5})$.

Proof. Note that

$$H(\operatorname{gen} L)=\{n\in\mathbb{N}_0 \mid n\equiv 0,1,4 \pmod 5\}.$$

Since L is a sublattice of a regular lattice $\langle 1,10\rangle$ (see Case II [$m=5$] (2)), L is regular. □

(4) $L=\langle 1,40\rangle$ is regular over $\mathbb{Q}(\sqrt{-5})$.

Proof. Note that

$$H(\operatorname{gen} L)=\{n\in\mathbb{N}_0 \mid n\equiv 0,1,4,5,6 \pmod 8\}\cap\{n\in\mathbb{N}_0 \mid n\equiv 0,1,4 \pmod 5\}.$$

Consider regular lattices $\langle 1,8\rangle$ and $\langle 1,10\rangle$ (see Case II [$m=5$] (1), (2)). Since $5\langle 1,8\rangle=\langle 5,40\rangle$ is a sublattice of L, L represents all positive integers $n\equiv 0,1,4,5,6 \pmod 8$ and $n\equiv 0 \pmod 5$. Since $4\langle 1,10\rangle=\langle 4,40\rangle$ is a sublattices of L, L represents all positive integers $n\equiv 0,1,4 \pmod 5$ and $n\equiv 0 \pmod 4$. On the other hand, the associated quadratic lattice of L is

$$x\overline{x}+40y\overline{y}=x_1^2+5x_2^2+40y_1^2+160y_2^2.$$

Since it has a regular sublattice $x_1^2+5x_2^2+40y_1^2$ [12], which represents all positive integers $n\equiv 1,5,6 \pmod 8$ and $n\equiv 1,4 \pmod 5$, these n are represented by L. Therefore L is regular. □

(5) $L=\begin{pmatrix}2 & -1+\omega\\ -1+\overline{\omega} & 3\end{pmatrix}\perp\langle 4\rangle$ is regular over $\mathbb{Q}(\sqrt{-5})$.

Proof. Note that

$$H(\operatorname{gen} L)=\{n\in\mathbb{N}_0 \mid n\equiv 0 \pmod 2 \text{ or } n\equiv 3 \pmod 4\}.$$

The associated quadratic lattice of L is

$$2x_1^2+2x_1x_2+3x_2^2+4y_1^2+20y_2^2.$$

Note that it contains regular sublattices $\ell_1 = 2x_1^2 + 2x_1x_2 + 3x_2^2 + 4y_1^2$ and $\ell_2 = 2x_1^2 + 2x_1x_2 + 3x_2^2 + 20y_2^2$ [12]. The lattice ℓ_1 represents all positive integers n if $n \equiv 0, 2, 3 \pmod 4$ and $n \neq 5^d u_+$. The lattice ℓ_2 represents all positive integers n if $n \equiv 0, 2, 3 \pmod 4$ and $n \neq 5^e u_+$. Hence $n \to L$ if $n \equiv 0 \pmod 2$ or $n \equiv 3 \pmod 4$. Therefore L is regular. ☐

(6) $L = \begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix} \perp \langle 5\rangle$ is regular over $\mathbb{Q}(\sqrt{-5})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 2, 3 \pmod 5\}.$$

Since $\langle 1\rangle \perp \begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix}$ is universal [10] and $5\left(\langle 1\rangle \perp \begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix}\right)$ is a sublattice of L, $n \to L$ for all $n \equiv 0 \pmod 5$. On the other hand, the associated quadratic lattice of L is

$$2x_1^2 + 2x_1x_2 + 3x_2^2 + 5y_1^2 + 25y_2^2.$$

Since it has a regular sublattice $2x_1^2 + 2x_1x_2 + 3x_2^2 + 5y_1^2$ [12], which represents all positive integers $n \equiv 2, 3 \pmod 5$, $n \to L$ for all positive integers $n \equiv 2, 3 \pmod 5$. Therefore L is regular. ☐

(7) $L = \begin{pmatrix} 2 & -1+\omega \\ -1+\overline{\omega} & 3 \end{pmatrix} \perp \langle 20\rangle$ is regular over $\mathbb{Q}(\sqrt{-5})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 2, 3 \pmod 4\} \cap \{n \in \mathbb{N}_0 \mid n \equiv 0, 2, 3 \pmod 5\}.$$

The associated quadratic lattice $\widetilde{L}$ of L is

$$2x_1^2 + 2x_1x_2 + 3x_2^2 + 20y_1^2 + 100y_2^2.$$

We mentioned that a regular lattice $2x_1^2+2x_1x_2+3x_2^2+4y^2$ represents n if $n \equiv 0, 2, 3 \pmod 4$ (see Case II [$m = 5$] (5)). Since $5\left(2x_1^2 + 2x_1x_2 + 3x_2^2 + 4y^2\right)$ is a sublattice of $\widetilde{L}$, $5n \to L$ if $5n \equiv 3 \pmod 4$ or $5n \equiv 0 \pmod 2$. Now consider another sublattice $2x_1^2 + 2x_1x_2 + 3x_2^2 + 20y_1^2$ (see Case II [$m = 5$] (5)). It represents n, if $n \equiv 0, 2, 3 \pmod 4$ and $n \equiv 2, 3 \pmod 5$. Hence the lattice L is regular. ☐

Case II [$m = 6$]

(1) $L = \langle 1, 3\rangle$ is regular over $\mathbb{Q}(\sqrt{-6})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 1 \pmod 3\}.$$

The associated quadratic lattice of L is

$$x\overline{x} + 3y\overline{y} = x_1^2 + 6x_2^2 + 3y_1^2 + 18y_2^2.$$

Since it has a regular sublattice $x_1^2 + 3y_1^2 + 6x_2^2$ [12], which represents all positive integers $n \equiv 0, 1 \pmod 3$, L is regular. ☐

(2) $L = \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix} \perp 3\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$ is regular over $\mathbb{Q}(\sqrt{-6})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 2 \pmod 3\}.$$

Since $3\left(\langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}\right)$ is a sublattice of L and $\langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$ is universal [10], L represents all positive integers $n \equiv 0 \pmod 3$. Suppose $n \equiv 2 \pmod 3$. Put $n =$

n_1+n_2 with $n_1 \to \langle 1\rangle$ and $n_2 \to \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$. Then $n_1 \equiv 0,1 \pmod 3$ and $n_2 \equiv 0,2 \pmod 3$. We have $n_1 \equiv 0 \pmod 3$ and $n_2 \equiv 2 \pmod 3$ from $n \equiv 2 \pmod 3$. Since $n_1 \to \langle 1,6\rangle_{\mathbb{Z}}$ and $n_1 \equiv 0 \pmod 3$, $n_1 \to 3\langle 2,3\rangle_{\mathbb{Z}}$. Note that $3\langle 2,3\rangle_{\mathbb{Z}}$ is an associated quadratic form of $3\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$. Therefore $n = n_1+n_2 \to 3\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix} \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 3 \end{pmatrix}$ and L is regular. ☐

Case II $[m=7]$

(1) $L=\langle 1,7\rangle$ is regular over $\mathbb{Q}(\sqrt{-7})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0,1,2,4 \pmod 7\}.$$

Since $\langle 7,7\rangle = 7\langle 1,1\rangle$ is a sublattice of L and $\langle 1,1\rangle$ is universal [7], $n \to L$ for all $n \equiv 0 \pmod 7$. On the other hand, the associated quadratic lattice of L is

$$x\overline{x} + 7y\overline{y} = x_1^2 + x_1x_2 + 2x_2^2 + 7y_1^2 + 7y_1y_2 + 14y_2^2.$$

Since it has a regular sublattice $x_1^2+x_1x_2+2x_2^2+7y_1^2$ [12], which represents all positive integers $n \equiv 1,2,4 \pmod 7$, $n \to L$ for all positive integers $n \equiv 1,2,4 \pmod 7$. Therefore L is regular. ☐

(2) $L=\langle 1,14\rangle$ is regular over $\mathbb{Q}(\sqrt{-7})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0,1,2,4 \pmod 7\}.$$

Since $7\langle 1,2\rangle = \langle 7,14\rangle$ is a sublattice of L and $\langle 1,2\rangle$ is universal [7], $n \to L$ for all positive integers $n \equiv 0 \pmod 7$. So we may assume n is not a multiple of 7 throughout this proof.

Let $\widetilde{L}$ be the associated quadratic lattice of L, then

$$\widetilde{L} = x\overline{x}+14y\overline{y} = x_1^2+x_1x_2+2x_2^2+14y_1^2+14y_1y_2+28y_2^2 = \begin{pmatrix} 1 & 1/2 \\ 1/2 & 2 \end{pmatrix}_{\mathbb{Z}} \perp \begin{pmatrix} 14 & 7 \\ 7 & 28 \end{pmatrix}_{\mathbb{Z}}.$$

Note that $\langle 1,7\rangle_{\mathbb{Z}}$ and $\langle 2,14\rangle_{\mathbb{Z}}$ are sublattices of $\begin{pmatrix} 1 & 1/2 \\ 1/2 & 2 \end{pmatrix}_{\mathbb{Z}}$.

(i) $n \equiv 0 \pmod 2$: Consider an integral ternary lattice $K = \langle 2\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}_{\mathbb{Z}}$ whose class number is one [1]. It represents all positive even integers not divisible by 7. Note that $\widetilde{L}$ has a sublattice $\langle 2,14\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 14 & 7 \\ 7 & 28 \end{pmatrix}_{\mathbb{Z}} = \langle 2\rangle_{\mathbb{Z}} \perp 7\left(\langle 2\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}_{\mathbb{Z}}\right) = \langle 2\rangle_{\mathbb{Z}} \perp K^7$. Suppose $n \geq 72$. Then $n \to \langle 2\rangle_{\mathbb{Z}} \perp K^7$ from the following identities.

$$n \equiv 1 \pmod 7 \Rightarrow n = 14k+8 = \begin{cases} 2\cdot 2^2 + 7\cdot 2k & \text{if } 7 \nmid k \\ 2\cdot 5^2 + 7\cdot 2(k-3) & \text{if } 7 \mid k \end{cases}$$

$$n \equiv 2 \pmod 7 \Rightarrow n = 14k+2 = \begin{cases} 2\cdot 1^2 + 7\cdot 2k & \text{if } 7 \nmid k \\ 2\cdot 6^2 + 7\cdot 2(k-5) & \text{if } 7 \mid k \end{cases}$$

$$n \equiv 4 \pmod 7 \Rightarrow n = 14k+4 = \begin{cases} 2\cdot 3^2 + 7\cdot 2(k-1) & \text{if } 7 \nmid (k-1) \\ 2\cdot 4^2 + 7\cdot 2(k-2) & \text{if } 7 \mid (k-1) \end{cases}$$

(ii) $n \equiv 1 \pmod 4$: Consider an integral ternary lattice $N = \langle 1\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}_{\mathbb{Z}}$. We know that the class number of N is two [1] and $\operatorname{gen} N$ represents all positive integers not divisible by 7. Since N_2 is isotropic over $\mathbb{Z}_2$, N represents positive integers $4k$ not divisible by 7 [9]. Note that $\widetilde{L}$ has a sublattice $\langle 1, 7\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 14 & 7 \\ 7 & 28 \end{pmatrix}_{\mathbb{Z}} = \langle 1\rangle_{\mathbb{Z}} \perp 7\left(\langle 1\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}_{\mathbb{Z}}\right) = \langle 1\rangle_{\mathbb{Z}} \perp N^7$. Suppose $n \geq 169$. Then $n \to \langle 1\rangle_{\mathbb{Z}} \perp N^7$ from the following identities.

$$n \equiv 1 \pmod 7 \Rightarrow n = 28k + 1 = \begin{cases} 1^2 + 7\cdot 4k & \text{if } 7 \nmid k \\ 13^2 + 7\cdot 4(k-6) & \text{if } 7 \mid k \end{cases}$$

$$n \equiv 2 \pmod 7 \Rightarrow n = 28k + 9 = \begin{cases} 3^2 + 7\cdot 4k & \text{if } 7 \nmid k \\ 11^2 + 7\cdot 4(k-4) & \text{if } 7 \mid k \end{cases}$$

$$n \equiv 4 \pmod 7 \Rightarrow n = 28k + 25 = \begin{cases} 5^2 + 7\cdot 4k & \text{if } 7 \nmid k \\ 9^2 + 7\cdot 4(k-2) & \text{if } 7 \mid k \end{cases}$$

(iii) $n \equiv 3 \pmod 8$: Consider an integral ternary lattice $M = \langle 11\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}_{\mathbb{Z}}$. We know that the class number of M is 5 [1] and $\operatorname{gen} M$ represents all positive integers not divisible by 7. Since M_2 is isotropic over $\mathbb{Z}_2$, M represents positive integers 4^4k not divisible by 7. Note that $\widetilde{L}$ has a sublattice $\langle 11, 77\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 14 & 7 \\ 7 & 28 \end{pmatrix}_{\mathbb{Z}} = \langle 11\rangle_{\mathbb{Z}} \perp 7\left(\langle 11\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}_{\mathbb{Z}}\right) = \langle 11\rangle_{\mathbb{Z}} \perp M^7$. There are $a \in \{1, 3, 5\}$ and $0 \leq b \leq 4^3 - 1$ such that

$$n \equiv 11a^2 \pmod 7 \text{ and } n \equiv 11(a + 14b)^2 \pmod{4^4}.$$

Let $k = \frac{n - 11(a+14b)^2}{7\cdot 4^4}$ and $\ell = \frac{n - 11(a+14b-7\cdot 2^7)^2}{7\cdot 4^4} = k - 11(a + 14b) + 11\cdot 7\cdot 2^6$. Then k and ℓ are positive integers if we assume that $n \geq 11(7\cdot 2^7)^2 = 8,830,976$. Note that both k and ℓ are not divisible by 7. Thus $n \to \langle 11\rangle_{\mathbb{Z}} \perp M^7$.

$$n = \begin{cases} 11(a + 14b)^2 + 7\cdot 4^4 k & \text{if } 7 \nmid k \\ 11(a + 14b - 7\cdot 2^7)^2 + 7\cdot 4^4 \ell & \text{if } 7 \mid k \end{cases}$$

(iv) $n \equiv 7 \pmod 8$: Consider an integral ternary lattice $R = \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}_{\mathbb{Z}} \perp \langle 23\rangle_{\mathbb{Z}}$. Then $\widetilde{L}$ contains $\langle 23, 161\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 14 & 7 \\ 7 & 28 \end{pmatrix}_{\mathbb{Z}} = \langle 23\rangle_{\mathbb{Z}} \perp R^7$. We know that the class number of R is 9 and the genus of R consists of 9 lattices [1]

$$R_1 = R = \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}_{\mathbb{Z}} \perp \langle 23\rangle_{\mathbb{Z}}, R_2 = \langle 1, 7, 23\rangle_{\mathbb{Z}}, R_3 = \begin{pmatrix} 4 & 1 \\ 1 & 6 \end{pmatrix}_{\mathbb{Z}} \perp \langle 7\rangle_{\mathbb{Z}},$$

$$R_4 = \begin{pmatrix} 2 & 1 \\ 1 & 12 \end{pmatrix}_{\mathbb{Z}} \perp \langle 7\rangle_{\mathbb{Z}}, R_5 = \begin{pmatrix} 3 & 1 \\ 1 & 8 \end{pmatrix}_{\mathbb{Z}} \perp \langle 7\rangle_{\mathbb{Z}}, R_6 = \begin{pmatrix} 3 & 1 & 1 \\ 1 & 4 & 0 \\ 1 & 0 & 15 \end{pmatrix}_{\mathbb{Z}},$$

$$R_7 = \langle 1\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 11 & 2 \\ 2 & 15 \end{pmatrix}_{\mathbb{Z}}, R_8 = \begin{pmatrix} 4 & 1 & 1 \\ 1 & 4 & 0 \\ 1 & 0 & 11 \end{pmatrix}_{\mathbb{Z}}, R_9 = \langle 1\rangle_{\mathbb{Z}} \perp \begin{pmatrix} 14 & 7 \\ 7 & 15 \end{pmatrix}_{\mathbb{Z}}.$$

Let f_{R_i} be the quadratic form corresponding to R_i. Then $f_R(x, y, z) = 2x^2 + 2xy + 4y^2 + 23z^2$. Note that $\operatorname{gen} R$ represents all positive integers not divisible by 7. If

$7 \nmid k$, then $4^7k \to R$ from the following identities:

$$\begin{aligned}
f_R(4y, -x-y, 2z) &= 4f_{R_2}(x,y,z),\\
f_R(8z, -4x-y-2z, 2y) &= 4^2 f_{R_3}(x,y,z),\\
f_R(16y, -2x-8y-4z, 4x) &= 4^3 f_{R_4}(x,y,z),\\
f_R(16y, 5x-4y-6z, 2x+4z) &= 4^3 f_{R_5}(x,y,z),\\
f_R(16y+32z, 10x+7y-20z, 4x-2y+8z) &= 4^4 f_{R_6}(x,y,z),\\
f_R(16x-16y+64z, 7x+33y-40z, -2x+18y+16z) &= 4^5 f_{R_7}(x,y,z),\\
f_R(48x-64y-32z, -47x-28y+66z, 18x+8y+36z) &= 4^6 f_{R_8}(x,y,z),\\
f_R(48x-240y-304z, -47x+123y-65z, 18x+70y+14z) &= 4^7 f_{R_9}(x,y,z).
\end{aligned}$$

There are $a \in \{1,3,5\}$ and $0 \le b \le 4^6 - 1 = 4095$ such that

$$n \equiv 23a^2 \pmod 7 \ \text{ and } n \equiv 23(a+14b)^2 \pmod{4^7}.$$

Let $k = \frac{n-23(a+14b)^2}{7\cdot 4^7}$ and $\ell = \frac{n-23(a+14b-7\cdot 4^7)^2}{7\cdot 4^7} = k - 23\cdot 2(a+14b) + 23\cdot 7\cdot 4^7$. Then k and ℓ are positive integers if we assume that $n \ge 23(7\cdot 2^{14})^2 = 302,526,758,912$. Note that both k and ℓ are not divisible by 7. Thus $n \to \langle 23\rangle_{\mathbb{Z}} \perp R^7$.

$$n = \begin{cases} 23(a+14b)^2 + 7\cdot 4^7 k & \text{if } 7 \nmid k \\ 23(a+14b-7\cdot 4^7)^2 + 7\cdot 4^7 \ell & \text{if } 7 \mid k \end{cases}$$

We checked over n represented by $\widetilde{L}$ for all $n \le 302,526,758,912$ by computer calculation. Therefore L is regular. □

Case II $[m = 10]$

(1) $L = \langle 1, 5\rangle$ is regular over $\mathbb{Q}(\sqrt{-10})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 1, 4 \pmod 5\}.$$

The associated quadratic lattice is

$$x\overline{x} + 5y\overline{y} = x_1^2 + 10x_2^2 + 5y_1^2 + 50y_2^2.$$

It has a regular sublattice $\ell = x_1^2 + 10x_2^2 + 5y_1^2$ [12], which represents all positive integers except $n = 5^e u_-$. Now suppose $n = 5^2(5k+2)$ or $n = 5^2(5k+3)$, for some integer k. Since $n - 50\cdot 1^2$ or $n - 50\cdot 2^2$ is represented by ℓ, hence $n \to L$ when $n \ge 50\cdot 2^2$. It is an easy work to check $n \to L$ for $n < 50\cdot 2^2$. Therefore L is regular. □

Case II $[m = 11]$

(1) $L = \langle 1, 4\rangle$ is regular over $\mathbb{Q}(\sqrt{-11})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 1 \pmod 2 \text{ or } n \equiv 0 \pmod 4\}.$$

Since L contains $4\langle 1,1\rangle = \langle 4,4\rangle$ and $\langle 1,1\rangle$ is universal [7], $n \to L$ for all positive integers $n \equiv 0 \pmod 4$. Note that the associated quadratic lattice of L is

$$x\overline{x} + 4y\overline{y} = x_1^2 + x_1x_2 + 3x_2^2 + 4y_1^2 + 4y_1y_2 + 12y_2^2.$$

Since $x_1^2 + x_1x_2 + 3x_2^2 + 4y_1^2$ is regular [12] and represents all positive integers $n \equiv 1 \pmod 2$ and $n \ne 11^d u_+$, these n are represented by L. Suppose $n \equiv 1 \pmod 2$ and $n = 11^d u_+$. Since $n/11 \to L$, $n \to L$. Therefore L is regular. □

(2) $L = \langle 1, 11\rangle$ is regular over $\mathbb{Q}(\sqrt{-11})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 1, 3, 4, 5, 9 \pmod{11}\}.$$

Since L contains $11\langle 1, 1\rangle = \langle 11, 11\rangle$ and $\langle 1, 1\rangle$ is universal [7], $n \to L$ for all positive integers $n \equiv 0 \pmod{11}$. Note that the associated quadratic lattice of L is

$$x\overline{x} + 11y\overline{y} = x_1^2 + x_1x_2 + 3x_2^2 + 11y_1^2 + 11y_1y_2 + 33y_2^2.$$

Since $x_1^2 + x_1x_2 + 3x_2^2 + 11y_1^2$ is regular [12] and represents all positive integers $n \equiv 1, 3, 4, 5, 9 \pmod{11}$, L represents these all integers. Therefore L is regular. ☐

(3) $L = \langle 1, 44\rangle$ is regular over $\mathbb{Q}(\sqrt{-11})$.

Proof. Note that

$$\begin{aligned} H(\operatorname{gen} L) = &\{n \in \mathbb{N}_0 \mid n \equiv 1 \pmod 2 \text{ or } n \equiv 0 \pmod 4\} \\ &\cap \{n \in \mathbb{N}_0 \mid n \equiv 0, 1, 3, 4, 5, 9 \pmod{11}\}. \end{aligned}$$

The associated quadratic lattice of L is

$$x_1^2 + x_1x_2 + 3x_2^2 + 44y_1^2 + 44y_1y_2 + 132y_2^2.$$

The sublattice $x_1^2 + x_1x_2 + 3x_2^2 + 44y_1^2$ is regular [12] and it represents all positive integers $n \equiv 0, 1, 3 \pmod 4$ and $n \neq 11^e u_-$. Since $11\langle 1, 4\rangle$ is a sublattice of L and $\langle 1, 4\rangle$ is regular (see Case II $[m = 11]$ (1)), L represents all positive integers $n \equiv 0 \pmod{11}$ and $n \equiv 0, 1, 3 \pmod 4$. Therefore L is regular. ☐

Case II $[m = 15]$

(1) $L = \langle 1, 3\rangle$ is regular over $\mathbb{Q}(\sqrt{-15})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 1 \pmod 3\}.$$

Since $3\left(\langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2\end{pmatrix}\right)$ is a sublattice of L and $\langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2\end{pmatrix}$ is universal [10], $n \to L$ for all positive integers $n \equiv 0 \pmod 3$. Suppose $n \equiv 1 \pmod 3$. Then $n \to \langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2\end{pmatrix}$ by universality. If $n = n_1 + n_2$ such that $n_1 \to \langle 1\rangle$ and $n_2 \to \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2\end{pmatrix}$, then we have $n_1 \equiv 1 \pmod 3$ and $n_2 \equiv 0 \pmod 3$, because $\langle 1\rangle$ represents $n_1 \equiv 0, 1 \pmod 3$ and $\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2\end{pmatrix}$ represents $n_2 \equiv 0, 2 \pmod 3$. Since $n_2 = 2x_1^2 + x_1x_2 + 2x_2^2$ has an integral solution with $x_1 + x_2 \equiv 0 \pmod 3$, $n_2 = 3\left[x_1 - (1+\omega)\frac{x_1+x_2}{3}\right]\overline{\left[x_1 - (1+\omega)\frac{x_1+x_2}{3}\right]} \to \langle 3\rangle$. So $n = n_1 + n_2 \to \langle 1, 3\rangle$ and L is regular. ☐

(2) $L = \langle 1, 5\rangle$ is regular over $\mathbb{Q}(\sqrt{-15})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 1, 4 \pmod 5\}.$$

The associated quadratic lattice of L is

$$x\overline{x} + 5y\overline{y} = x_1^2 + x_1x_2 + 4x_2^2 + 5y_1^2 + 5y_1y_2 + 20y_2^2.$$

Since it has a regular sublattice $x_1^2 + x_1x_2 + 4x_2^2 + 5y_1^2$ [12], which represents all positive integers $n \equiv 1, 4 \pmod 5$, L represents all positive integers $n \equiv 1, 4 \pmod 5$. Since $5\left(\langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2\end{pmatrix}\right)$ is a sublattice of L and $\langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2\end{pmatrix}$ is universal [10], $n \to L$ for all positive integers $n \equiv 0 \pmod 5$. Therefore L is regular. ☐

(3) $L = \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix} \perp \langle 5 \rangle$ is regular over $\mathbb{Q}(\sqrt{-15})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 2, 3 \pmod 5\}.$$

The associated quadratic lattice $\widetilde{L}$ of L

$$2x_1^2 + x_1x_2 + 2x_2^2 + 5y_1^2 + 5y_1y_2 + 20y_2^2$$

has a regular sublattice $2x_1^2 + x_1x_2 + 2x_2^2 + 5y^2$ [12], which represents all positive integers $n \equiv 0, 2, 3 \pmod 5$ and $n \neq 3^d u_-$. Suppose $n \equiv 0, 2, 3 \pmod 5$ and $n = 3^d u_-$. We may consider $n = 3(15k+5)$, $n = 3(15k+11)$ or $n = 3(15k+14)$ for some nonnegative integers k. Consider a quadratic sublattice $2x_1^2+x_1x_2+2x_2^2+5y^2+75z^2$ of $\widetilde{L}$. Then $n - 75 \cdot 1^2$ is not of the form $3^d u_-$ and hence it is represented by $2x_1^2 + x_1x_2 + 2x_2^2 + 5y^2$. The representability of $3 \cdot 5$, $3 \cdot 20$, $3 \cdot 11$, $3 \cdot 14$ by L is easily checked. Therefore, L is regular. ☐

(4) $L = \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix} \perp 3 \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$ is regular over $\mathbb{Q}(\sqrt{-15})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 2 \pmod 3\}.$$

Since $3 \to \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$, $3\left(\langle 1 \rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}\right)$ is a sublattice of L. Since the lattice $\langle 1 \rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$ is universal [10], L represents all positive integers $n \equiv 0 \pmod 3$. Suppose $n \equiv 2 \pmod 3$. If $n = n_1 + n_2$ such that $n_1 \to \langle 1 \rangle$ and $n_2 \to \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$, then we have $n_1 \equiv 0, 1 \pmod 3$ and $n_2 \equiv 0, 2 \pmod 3$. Since $n \equiv 2 \pmod 3$, $n_1 \equiv 0 \pmod 3$. Then $n_1 = (x_1 + x_2\omega)\overline{(x_1 + x_2\omega)}$ has an integral solution with $x_1 \equiv x_2 \pmod 3$. Since $n_1 = 6\alpha\overline{\alpha} + 3\omega\alpha\overline{\beta} + 3\overline{\omega}\overline{\alpha}\beta + 6\beta\overline{\beta}$ with $\alpha = \frac{x_1 - y_1}{3}$ and $\beta = -\frac{x_1 + 2x_2}{3}$, $n_1 \to 3\begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$ and L is regular. ☐

(5) $L = \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix} \perp \langle 9 \rangle$ is regular over $\mathbb{Q}(\sqrt{-15})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 2 \pmod 3 \text{ or } n \equiv 0, 3 \pmod 9\}.$$

Since $3\langle 1, 3 \rangle$ is a sublattice of L and $\langle 1, 3 \rangle$ is a regular lattice which represents all positive integers $n \equiv 0, 1 \pmod 3$ (see Case$[m = 15]$ (1)), L represents all positive integers $n \equiv 0, 3 \pmod 9$. Suppose $n \equiv 2 \pmod 3$. The associated quadratic lattice $\widetilde{L}$ of L

$$x_1^2 + x_1x_2 + x_2^2 + 9y_1^2 + 9y_1y_2 + 36y_2^2$$

has a regular sublattice $\ell = x_1^2 + x_1x_2 + x_2^2 + 9y_1^2$ [12] and it represents all positive integers n if $n \neq 5^d u_-$. Suppose $n = 5u_-$. Then $n = 5(15k+7), 5(15k+13), 5(45k+33)$ or $5(45k+42)$. The quadratic lattice $\widetilde{L}$ has the sublattice $x_1^2+x_1x_2+x_2^2+9y^2+135z^2$. Then, $5(15k+7) - 135 \cdot 2^2$, $5(15k+13) - 135 \cdot 1^2$, $5(45k+33) - 135 \cdot 1^2$ and $5(45k+42) - 135 \cdot 2^2$ are all represented by ℓ when $n \geq 135 \cdot 2^2$. The representability for $n < 135 \cdot 2^2$ is easily checked. Therefore L is regular. ☐

(6) $L = \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix} \perp \langle 15 \rangle$ is regular over $\mathbb{Q}(\sqrt{-15})$.

Proof. Note that

$$H(\operatorname{gen} L) = \{n \in \mathbb{N}_0 \mid n \equiv 0, 2 \pmod 3 \text{ and } n \equiv 0, 2, 3 \pmod 5\}.$$

Since $15\left(\langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}\right)$ is a sublattice of L and $\langle 1\rangle \perp \begin{pmatrix} 2 & \omega \\ \overline{\omega} & 2 \end{pmatrix}$ is universal, L represents all positive integers divisible by 15. The associated quadratic lattice $\widetilde{L}$ of L is

$$2x_1^2 + x_1x_2 + 2x_2^2 + 15y_1^2 + 15y_1y_2 + 60y_2^2.$$

The sublattice $\ell = 2x_1^2 + x_1x_2 + 2x_2^2 + 15y_1^2$ is regular [12] and it represents all positive integers $n \equiv 0, 2 \pmod 3$ and $n \neq 5^{2s}u_+$ with $s \geq 1$.

It is enough to consider positive integers n such that $n \equiv 2 \pmod 3$ and $n = 5^2u_+$. Suppose $n = 5^2(15k + 11)$ or $n = 5^2(15k + 14)$. The quadratic lattice $\widetilde{L}$ has a sublattice $2x_1^2 + x_1x_2 + 2x_2^2 + 15y_1^2 + 225z^2$. Then, $n - 225 \cdot 1^2$ and $n - 225 \cdot 2^2$ are represented by ℓ when $n \geq 225 \cdot 2^2$. The representability for $n < 225 \cdot 2^2$ is easily checked. Therefore L is regular. □

Theorem 1. *There are* 68 *binary regular normal Hermitian lattices, including* 9 *nondiagonal lattices, up to isometry over* $\mathbb{Q}(\sqrt{-m})$ *with positive square-free integers* m*. In Table 13,* 25 *binary universal Hermitian lattices are marked with* †.

$\mathbb{Q}(\sqrt{-m})$	binary regular Hermitian lattices (†: universal)
$\mathbb{Q}(\sqrt{-1})$	$\langle 1,1\rangle^\dagger$, $\langle 1,2\rangle^\dagger$, $\langle 1,3\rangle^\dagger$, $\langle 1,4\rangle$, $\langle 1,8\rangle$, $\langle 1,16\rangle$, $\begin{pmatrix} 2 & -1+\omega_1 \\ -1+\overline{\omega}_1 & 3 \end{pmatrix}$, $\begin{pmatrix} 3 & -1+\omega_1 \\ -1+\overline{\omega}_1 & 6 \end{pmatrix}$, $\begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$
$\mathbb{Q}(\sqrt{-2})$	$\langle 1,1\rangle^\dagger$, $\langle 1,2\rangle^\dagger$, $\langle 1,3\rangle^\dagger$, $\langle 1,4\rangle^\dagger$, $\langle 1,5\rangle^\dagger$, $\langle 1,8\rangle$, $\langle 1,16\rangle$, $\langle 1,32\rangle$, $\begin{pmatrix} 2 & \omega_2 \\ \overline{\omega}_2 & 5 \end{pmatrix}$
$\mathbb{Q}(\sqrt{-3})$	$\langle 1,1\rangle^\dagger$, $\langle 1,2\rangle^\dagger$, $\langle 1,3\rangle$, $\langle 1,4\rangle$, $\langle 1,6\rangle$, $\langle 1,9\rangle$,$\langle 1,12\rangle$, $\langle 1,36\rangle$, $\langle 2,3\rangle$, $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, $\begin{pmatrix} 2 & 1 \\ 1 & 5 \end{pmatrix}$, $\begin{pmatrix} 3 & 1+\omega_3 \\ 1+\overline{\omega}_3 & 5 \end{pmatrix}$, $\begin{pmatrix} 5 & 2 \\ 2 & 8 \end{pmatrix}$
$\mathbb{Q}(\sqrt{-5})$	$\langle 1,2\rangle^\dagger$, $\langle 1\rangle \perp \begin{pmatrix} 2 & -1+\omega_5 \\ -1+\overline{\omega}_5 & 3 \end{pmatrix}^\dagger$, $\langle 1,8\rangle$, $\langle 1,10\rangle$, $\langle 1,40\rangle$, $\langle 1\rangle \perp 5\begin{pmatrix} 2 & -1+\omega_5 \\ -1+\overline{\omega}_5 & 3 \end{pmatrix}$, $\begin{pmatrix} 2 & -1+\omega_5 \\ -1+\overline{\omega}_5 & 3 \end{pmatrix} \perp \langle 4\rangle$, $\begin{pmatrix} 2 & -1+\omega_5 \\ -1+\overline{\omega}_5 & 3 \end{pmatrix} \perp \langle 5\rangle$, $\begin{pmatrix} 2 & -1+\omega_5 \\ -1+\overline{\omega}_5 & 3 \end{pmatrix} \perp \langle 20\rangle$
$\mathbb{Q}(\sqrt{-6})$	$\langle 1\rangle \perp \begin{pmatrix} 2 & \omega_6 \\ \overline{\omega}_6 & 3 \end{pmatrix}^\dagger$, $\langle 1,3\rangle$, $\begin{pmatrix} 2 & \omega_6 \\ \overline{\omega}_6 & 3 \end{pmatrix} \perp 3\begin{pmatrix} 2 & \omega_6 \\ \overline{\omega}_6 & 3 \end{pmatrix}$
$\mathbb{Q}(\sqrt{-7})$	$\langle 1,1\rangle^\dagger$, $\langle 1,2\rangle^\dagger$, $\langle 1,3\rangle^\dagger$, $\langle 1,7\rangle$, $\langle 1,14\rangle$, $\begin{pmatrix} 3 & \omega_7 \\ \overline{\omega}_7 & 3 \end{pmatrix}$
$\mathbb{Q}(\sqrt{-10})$	$\langle 1\rangle \perp \begin{pmatrix} 2 & \omega_{10} \\ \overline{\omega}_{10} & 5 \end{pmatrix}^\dagger$, $\langle 1,5\rangle$
$\mathbb{Q}(\sqrt{-11})$	$\langle 1,1\rangle^\dagger$, $\langle 1,2\rangle^\dagger$, $\langle 1,4\rangle$, $\langle 1,11\rangle$, $\langle 1,44\rangle$
$\mathbb{Q}(\sqrt{-15})$	$\langle 1\rangle \perp \begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix}^\dagger$, $\langle 1,3\rangle$,$\langle 1,5\rangle$, $\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix} \perp \langle 5\rangle$, $\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix} \perp 3\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix}$, $\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix} \perp \langle 9\rangle$, $\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix} \perp \langle 15\rangle$
$\mathbb{Q}(\sqrt{-19})$	$\langle 1,2\rangle^\dagger$
$\mathbb{Q}(\sqrt{-23})$	$\langle 1\rangle \perp \begin{pmatrix} 2 & \omega_{23} \\ \overline{\omega}_{23} & 3 \end{pmatrix}^\dagger$, $\langle 1\rangle \perp \begin{pmatrix} 2 & -1+\omega_{23} \\ -1+\overline{\omega}_{23} & 3 \end{pmatrix}^\dagger$
$\mathbb{Q}(\sqrt{-31})$	$\langle 1\rangle \perp \begin{pmatrix} 2 & \omega_{31} \\ \overline{\omega}_{31} & 4 \end{pmatrix}^\dagger$, $\langle 1\rangle \perp \begin{pmatrix} 2 & -1+\omega_{31} \\ -1+\overline{\omega}_{31} & 4 \end{pmatrix}^\dagger$

TABLE 13. Binary regular Hermitian lattices: normal case

Remark 1. *Rokicki listed* 55 *binary regular* diagonal *Hermitian normal lattices over* $\mathbb{Q}(\sqrt{-m})$ *including a candidate* $\langle 1, 14\rangle$ *over* $\mathbb{Q}(\sqrt{-7})$ *up to isometry. We confirm her list and make up a list of all binary regular Hermitian normal lattices* *including (1) four diagonal lattices* $\langle 1\rangle \perp 5\begin{pmatrix} 2 & -1+\omega_5 \\ -1+\overline{\omega}_5 & 3 \end{pmatrix}$ *over* $\mathbb{Q}(\sqrt{-5})$, $\begin{pmatrix} 2 & \omega_6 \\ \overline{\omega}_6 & 3 \end{pmatrix} \perp 3\begin{pmatrix} 2 & \omega_6 \\ \overline{\omega}_6 & 3 \end{pmatrix}$ *over* $\mathbb{Q}(\sqrt{-6})$*, and* $\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix} \perp \langle 5\rangle$, $\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix} \perp \langle 9\rangle$ *over* $\mathbb{Q}(\sqrt{-15})$ *which she missed, (2) a universal lattice* $\langle 1, 3\rangle$ *over* $\mathbb{Q}(\sqrt{-7})$ *uncaught at that time and (3) nine nondiagonal lattices. Also we provide the complete proof of regularity of each lattice. Some of Rokicki's proofs are corrected and simplified.*

Remark 2. *Among the Hermitian lattices, we have some interesting lattices. For example, a lattice* $\begin{pmatrix} 2 & \omega_6 \\ \overline{\omega}_6 & 3 \end{pmatrix} \perp 3\begin{pmatrix} 2 & \omega_6 \\ \overline{\omega}_6 & 3 \end{pmatrix}$ *is isometric to* $\begin{pmatrix} 9 & 4\omega_6 \\ 4\overline{\omega}_6 & 11 \end{pmatrix}$ *over* $\mathbb{Q}(\sqrt{-6})$*. The former is a direct sum of nonfree unary lattices and the latter is a binary free lattice. And we have another example of the same phenomenon. A lattice* $\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix} \perp 3\begin{pmatrix} 2 & \omega_{15} \\ \overline{\omega}_{15} & 2 \end{pmatrix}$ *is isometric to* $\begin{pmatrix} 8 & -3+4\omega_{15} \\ -3+4\overline{\omega}_{15} & 8 \end{pmatrix}$ *over* $\mathbb{Q}(\sqrt{-15})$*. On the other hand, although both* $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ *over* $\mathbb{Q}(\sqrt{-3})$ *and* $\begin{pmatrix} 8 & -3+4\omega_{15} \\ -3+4\overline{\omega}_{15} & 8 \end{pmatrix}$ *over* $\mathbb{Q}(\sqrt{-15})$ *have even diagonal entries, they can represent odd integers. So they are normal lattices.*

Remark 3. *The binary subnormal regular Hermitian lattices will be investigated in our next articles. Binary subnormal regular Hermitian lattices over* $\mathbb{Q}(\sqrt{-m})$ *with norm ideal* $2\mathfrak{o}$ *occur only when*

$$m = 1, 2, 5, 6, 10, 13, 14, 22, 29, 34, 37 \textit{ and } 38.$$

Also, we know that binary subnormal regular Hermitian lattices exist over $\mathbb{Q}(\sqrt{-m})$ *whose norm ideal is* $m\mathfrak{o}$*. For example,* $\begin{pmatrix} 3 & \sqrt{-3} \\ -\sqrt{-3} & 3 \end{pmatrix}$ *over* $\mathbb{Q}(\sqrt{-3})$ *is a binary subnormal regular Hermitian lattice with norm ideal* $3\mathfrak{o}$*. It is an impossible phenomenon for quadratic lattices over* $\mathbb{Z}$*.*

REFERENCES

[1] Brandt, H., Intrau, O.: *Tabellen reduzierter positiver ternärer quadratischer Formen*, Abh. Sachs. Akad. Wiss. Math.-Nat. Kl. **45** (1958)

[2] Chan, W. K., Earnest, A. G., Icaza, M. I., Kim, J. Y.: *Finiteness results for regular definite ternary quadratic forms over* $\mathbb{Q}(\sqrt{5})$, Inter. J. Number Theory, **3**, 541–556 (2007)

[3] Chan, W. K., Rokicki, A.: *Positive definite binary hermitian forms with finitely many exceptions*, J. Number Theory **124**, 167–180 (2007)

[4] Conway, J. H.: *Universal quadratic forms and the fifteen theorem*, Contemp. Math. **272**, 23–26 (2000)

[5] Dickson, L. E.: *Ternary quadratic forms and congruences*, Ann. of Math. **28**, 331–341 (1927)

[6] Earnest, A. G.: *An application of character sum inequalities to quaratic forms*, Number Theory (Karl Dilcher, ed.), CMS Conf. Proc. **15**, 155–158 (1994), Amer. Math. Soc., Providence, RI (1995)

[7] Earnest, A. G., Khosravani, A.: *Universal binary Hermitian forms*, Math. Comp. **66**, 1161–1168 (1997)

[8] Earnest, A. G., Khosravani, A.: *Representation of integers by positive definite binary Hermitian lattices over imaginary quadratic fields*, J. Number Theory **62**(2), 368–374 (1997)

[9] Icaza, M. I.: *Sums of squares of integral linear forms*, Acta Arith. **74**, 231–240 (1996)

[10] Iwabuchi, H.: *Universal binary positive definite Hermitian lattices*, Rocky Mountain J. Math. **30**(3), 951–959 (2000)

[11] Jacobson, N.: *A note on hermitian forms*, Amer. Math. Soc. **46**, 264–268 (1940)

[12] Jagy, W. C., Kaplansky, I., Schiemann, A.: *There are 913 Regular Ternary Forms*, Mathematika **44**, 332–341 (1997)

[13] Jones, B. W.: *The regularity of a genus of positive ternary quadratic forms*, Trans. Amer. Math. Soc. **33**(1), 111–124 (1931)

[14] Jones, B. W., Pall, G.: *Regular and semi-regular positive ternary quadratic forms*, Acta Math. **70**, 165–191 (1939)

[15] Kim, B. M.: *Complete determination of regular positive diagonal quaternary integral quadratic forms*, preprint

[16] Kim, B. M., Kim, J. Y., Park, P.-S.: *The fifteen theorem for universal Hermitian lattices over imaginary quadratic fields*, arXiv:0710.4991v1.

[17] Kim, J.-H., Park, P.-S.: *A few uncaught universal Hermitian forms*, Proc. Amer. Math. Soc. **135**, 47–49 (2007)

[18] Kim, M.-H., Park, P.-S.: *Ternary 2-universal Hermitian lattices over imaginary quadratic fields*, preprint

[19] O'Meara, O. T.: *Introduction to Quadratic Forms*, Spinger-Verlag, New York (1973)

[20] Otremba, G.: *Zur Theorie der hermiteschen Formen in imaginär-quadratischen Zahlkörpern*, J. Reine Angew. Math. **249**, 1–19 (1971)

[21] Rokicki, A.:*Finiteness results for definite n-regular and almost n-regular hermitian forms*, Ph.D. Thesis, Wesleyan University (2005)

[22] Watson, G. L.: *Some problems in the theory of numbers*, Ph.D. thesis, University of London (1953)

[23] Watson, G. L.: *The representation of integers by positive ternary quadratic forms*, Mathematika **1**, 104–110 (1954)

[24] Watson, G. L.: *Transformations of a quadratic form which do not increase the class-number*, Proc. London Math. Soc. (3) **12**, 577–587 (1962)

Department of Mathematics, Kangnung National University, Kangnung, 210-702, Korea

E-mail address: `kbm@kangnung.ac.kr`

Department of Mathematical Sciences, Seoul National University, san 56-1 Shinrim-dong Gwanak-gu, Seoul, 151-747, Korea

E-mail address: `jykim@math.snu.ac.kr`

School of Computational Sciences, Korea Institute for Advanced Study, Hoegiro 87, Dongdaemun-gu, Seoul, 130-722, Korea

E-mail address: `sung@kias.re.kr`